



Illinois Air Quality Report



2020

ILLINOIS ANNUAL AIR QUALITY REPORT 2020

**Illinois Environmental Protection Agency
Bureau of Air
1021 North Grand Avenue, East
P.O. Box 19276
Springfield, IL 62794-9276**

Printed on recycled paper

For additional information on air pollution, please see the Illinois EPA website,
<http://www.epa.illinois.gov/>, or write to:

Illinois Environmental Protection Agency
Bureau of Air
1021 N. Grand Ave., East
PO Box 19276
Springfield, IL 62794-9276

Acknowledgements

This document is produced by the Illinois Environmental Protection Agency; John Kim, Director.

Illinois EPA Bureau of Air personnel contributed their time and expertise to the development of this publication.

Illinois Annual Air Quality Report 2020

Contents

Tables	5
Figures	6
Executive Summary	8
Section 1: Air Pollutants: Sources, Health & Welfare Effects	10
Section 2: Statewide Summary of Air Quality	16
Section 3: Air Quality Index	21
Section 4: Statewide Summary of Point Source Emissions	28

Appendices

Appendix A: Air Sampling Network	34-46
Sampling Schedule	35
Distribution of Air Monitoring Equipment	38
Statewide Air Monitoring Locations	39
Appendix B: Air Quality Data Summary Tables	47-87
Air Quality Data Interpretation	47
Ozone Data	49
Particulate Matter (PM _{2.5}) Data	57
Particulate Matter (PM ₁₀) Data	65
Carbon Monoxide Data	69
Sulfur Dioxide Data	73
Nitrogen Dioxide Data	77
Lead Data	82
Filter Analysis Data	85
Toxic Compounds Data	87
Appendix C: Point Source Emission Inventory Summary Tables	87
Appendix D: Website Links	97

Tables

Table 1: Summary of National and Illinois Ambient Air Quality Standards.....	15
Table 2: Illinois Air Pollution Episode Levels.....	15
Table 3: Air Quality Index Categories	22
Table 4: Air Quality Index Health Concerns	22
Table 5: Air Quality Index Sectors in Illinois.....	23
Table 6: Distribution of Volatile Organic Material Emissions	29
Table 7: Distribution of Particulate Matter Emissions.....	30
Table 8: Distribution of Carbon Monoxide Emissions	31
Table 9: Distribution of Sulfur Dioxide Emissions.....	32
Table 10: Distribution of Nitrogen Oxide Emissions.....	33
Table A1: Non-Continuous Sampling Schedule	35
Table A2: Distribution of Air Monitoring Equipment.....	38
Table A3: Site Directory	40
Table A4: Monitoring Directory	43
Table B1: 1-Hour Ozone Exceedances	50
Table B2: 8-Hour Ozone Exceedances	51
Table B3: Ozone Highs	53
Table B4: Ozone Design Values	55
Table B5: PM _{2.5} 24-Hour Exceedances.....	58
Table B6: PM _{2.5} Highs	59
Table B7: PM _{2.5} 24-Hour Design Values.....	61
Table B8: PM _{2.5} Annual Design Values.....	63
Table B9: PM ₁₀ 24-Hour Exceedances.....	66
Table B10: PM ₁₀ 24-Hour Highs and Design Values	67
Table B11: PM ₁₀ Annual Design Values	68
Table B12: Carbon Monoxide Exceedances	70
Table B13: Carbon Monoxide Highs	71
Table B14: Carbon Monoxide 1-Hour and 8-Hour Design Values.....	72
Table B15: Sulfur Dioxide Exceedances	74
Table B16: Sulfur Dioxide Highs.....	75
Table B17: Sulfur Dioxide 1-Hour Design Values	76
Table B18: Nitrogen Dioxide 1-Hour Exceedances.....	78
Table B19: Nitrogen Dioxide Highs	79
Table B20: Nitrogen Dioxide 1-Hour Design Values.....	80
Table B21: Nitrogen Dioxide Annual Design Values.....	81
Table B22: Lead Highs	83
Table B23: Lead Design Values.....	84
Table B24: Filter Analysis Data.....	85
Table B25: Toxic Compounds	86
Table C1: Carbon Monoxide Point Source Emission Distribution	87
Table C2: Nitrogen Oxides Point Source Emission Distribution.....	88
Table C3: PM ₁₀ Point Source Emission Distribution.....	89
Table C4: Sulfur Dioxide Point Source Emission Distribution	90
Table C5: Volatile Organic Material Point Source Emission Distribution.....	91
Table C6: Estimated County Stationary Point Source Emissions.....	93
Table C7: Annual Estimated Emissions Trends.....	95
Table C8: Annual Source Reported Emissions Trends.....	96

Figures

Figure 1: Average 1-Hour Ozone Maximum	16
Figure 2: Average 8-Hour Ozone 4th High.....	16
Figure 3: Particulate Matter (PM _{2.5}) Annual Trends.....	17
Figure 4: Particulate Matter (PM ₁₀) 24-hour Trends.....	17
Figure 5: Carbon Monoxide Trends	18
Figure 6: Sulfur Dioxide 24-hour Trends.....	18
Figure 7: Nitrogen Dioxide Annual Trend.....	19
Figure 8: Lead Rolling 3-Month Maximum Trend	19
Figure 9: Air Quality Index Summaries by Sector.....	25
Figure 10: Estimated Volatile Organic Material Emissions Trend	29
Figure 11: Estimated Particulate Emissions Trend	30
Figure 12: Estimated Carbon Monoxide Emissions Trend	31
Figure 13: Estimated Sulfur Dioxide Emissions Trend.....	32
Figure 14: Estimated Nitrogen Oxide Emissions Trend	33

Executive Summary

This report presents a summary of air quality data collected throughout the State of Illinois during calendar year 2020. Data is presented for the six criteria pollutants (those for which air quality standards have been developed – particulate matter (PM₁₀ and PM_{2.5}), ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide, and lead – along with some heavy metals, volatile organic compounds and toxic compounds. Monitoring was conducted at 64 different site locations collecting data from 144 instruments.

In terms of the Air Quality Index (AQI) air quality during 2020 was either good or moderate 96% of the time throughout Illinois. There were five days when air quality was considered unhealthy (category red). This compares with three unhealthy days in 2019. The unhealthy days were due to elevated ozone and particulate concentrations in June and July. There were 21 days (17 for ozone, two for fine particulates, and two for a combination of fine particulates and ozone) when air quality in some part of Illinois was considered Unhealthy for Sensitive Groups (category orange). This compares with 13 Unhealthy for Sensitive Groups days reported in 2019. Air quality trends for most of the criteria pollutants are continuing to show downward or stable trends below the level of the standards.

Stationary point source emission data has again been included. The data in the report reflects information contained in Illinois EPA's Integrated Comprehensive Environmental Management System (ICEMAN) as of December 31, 2020. Emission estimates are for the calendar year 2020 and are for the pollutants: particulate matter, volatile organic material, sulfur dioxide, nitrogen oxides, and carbon monoxide. Emission trends of these pollutants have been given for the years 1998 to the present. Emissions reported with the Annual Emissions Report have been provided starting with 1998 and are currently available through 2019. There has been a trend toward decreasing emissions over this time period.

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

Ozone (O₃)

Photochemical oxidants result from a complex series of atmospheric reactions initiated by sunlight. When reactive (non-methane) hydrocarbons and nitrogen oxides accumulate in the atmosphere and are exposed to the ultraviolet component of sunlight, the formation of new compounds, including ozone and peroxyacetylnitrate, takes place.

Absorption of ultraviolet light energy by nitrogen dioxide results in its dissociation into nitric oxide and an oxygen atom. The oxygen atoms, for the most part, react with atmospheric molecular oxygen (O₂) to form ozone (O₃). In general, nitric oxide will react with ozone to re-form nitrogen dioxide, completing the cycle. A build-up of ozone above the equilibrium concentration, which is defined by the reaction cycle, results when nitrogen oxide reacts with non-methane hydrocarbons. Oxygen atoms from the hydrocarbon radical oxidize nitric oxide to nitrogen dioxide without ozone being used up. Thus, ozone concentrations are not depleted and can build up quickly.

Ozone can also be formed naturally in the atmosphere by electrical discharge and in the stratosphere by solar radiation. The former process is not capable of producing significant urban concentrations of this pollutant; however, there is some belief that incursion of ozone from the stratosphere can contribute significantly to elevated ground level concentrations of ozone under certain meteorological conditions.

Injury to vegetation is one of the earliest manifestations of photochemical air pollution, and sensitive plants are useful biological indicators of this type of pollution. The visible symptoms of photochemical oxidant produced injury to plants may be classified as:

- Acute injury, identified by cell collapse with subsequent development of necrotic patterns.
- Chronic injury, identified by necrotic patterns or with other pigmented patterns.

- Physiological effects, identified by growth alterations, reduced yields, and changes in the quality of plant products. The acute symptoms are generally characteristic of a specific photochemical oxidant, though chronic injury patterns are not. Ozone injury to leaves is identified as a striping or flecking. Adverse effects on sensitive vegetation have been observed from exposure to photochemical oxidant concentrations of about 100 micrograms per cubic meter (0.05 parts per million) for 4 hours.

Adverse effects on materials (rubber products and fabrics) from exposure to photochemical oxidants have not been precisely quantified, but have been observed at the levels presently occurring in many urban atmospheres.

Ozone accelerates the aging of many materials, resulting in rubber cracking, dye fading, and paint erosion. These effects are linearly related to the total dose of ozone and can occur at very low levels, given long duration exposures.

Ozone is a pulmonary irritant that affects the respiratory mucous membranes, other lung tissues, and respiratory functions. Clinical and epidemiological studies have demonstrated that ozone impairs the normal mechanical function of the lung, causing alterations in respiration – the most characteristic of which are shallow, rapid breathing and a decrease in pulmonary compliance. Exposure to ozone results in clinical symptoms such as chest tightness, coughing, and wheezing. Alterations in airway resistance can occur, especially to those with respiratory diseases (asthma, bronchitis, emphysema). These effects may occur in sensitive individuals, as well as in healthy exercising persons, at short-term ozone concentrations between 0.15 and 0.25 ppm.

Ozone exposure increases the sensitivity of the lung to bronchoconstrictive agents such as histamine, acetylcholine, and allergens, as well as increasing the individual's susceptibility to bacterial infection. Simultaneous exposure to ozone and sulfur dioxide can produce larger changes in pulmonary function than exposure to either pollutant alone.

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

Peroxyacetylnitrate (PAN) is an eye irritant, and its effects often occur in conjunction with the effects of ozone.

Two characteristics of ozone and photochemical oxidant exposures should be cited:

- Ozone itself is a primary cause of most of the health effects reported in toxicological and experimental human studies and the evidence for attributing many health effects to this substance alone is very compelling.
- Atmospheric photochemical substances are known to produce health effects, some of which are not attributable to pure ozone but may be caused by other photochemical substances in combination with ozone.

Particulate Matter (PM)

Not all air pollutants are in the gaseous form. Small solid particles and liquid droplets, collectively called particulates or aerosols, are also present in the air in great numbers and may constitute a pollution problem. Particulates entering the atmosphere differ in size and chemical composition. The effects of particulates on health and welfare are directly related to their size and chemical composition.

Particulate matter in the atmosphere consists of solids, liquids, and liquids-solids in combination. Suspended particulates generally refer to particles less than 100 micrometers in diameter (human hair is typically 100 micrometers thick). Particles larger than 100 micrometers will settle out of the air under the influence of gravity in a short period of time.

Typical sources emitting particles into the atmosphere are combustion of fossil fuels (ash and soot), industrial processes (metals, fibers, etc.), fugitive dust (wind and mechanical erosion of local soil), and photochemically produced particles (complex chain reactions between sunlight and gaseous pollutants). Combustion and photochemical products tend to be smaller in size (less than 1 micrometer);

fugitive dust and industrial products are typically larger in size (greater than 1 micrometer).

Particles which cause the most health and visibility difficulties are those less than 1.0 micrometer in size. These particles are also the most difficult to reduce in numbers by the various industrial removal techniques. Rainfall accounts for the major removal of these smaller particles from the air.

One of the major problems associated with high concentrations of particulates is that the interaction between the particles, sunlight, and atmospheric moisture can potentially result in the climatic effects and diminished visibility (haze). Particles play a key role in the formation of clouds, and emissions of large numbers of particles can, in some instances, result in local increases in cloud formation and, possibly, precipitation. Particles in the size range of 0.1 to 1.0 micrometers are the most efficient in scattering visible light (wavelength 0.4 to 0.7 micrometers) thereby reducing visibility. Particles combined with high humidity can result in the formation of haze which can cause hazardous conditions for the operation of motor vehicles and aircraft.

Particulate pollutants enter the human body by way of the respiratory system and their most immediate effects are upon this system. The size of the particle determines its depth of penetration into the respiratory system. Particles over 5 micrometers are generally deposited in the nose and throat. Those that do penetrate deeper in the respiratory system to the air ducts (bronchi) are often removed by ciliary action. Particles ranging in size from 0.5 - 5.0 micrometers in diameter can be deposited in the bronchi, with few reaching the air sacs (alveoli). Most particles deposited in the bronchi are removed by the cilia within hours. Particles less than 0.5 micrometer in diameter reach and may settle in the alveoli. The removal of particles from the alveoli is much less rapid and complete than from the larger passages. Some of the particles retained in the alveoli are absorbed into the blood.

Besides particulate size, the oxidation state, chemical composition, concentration, and length of time in the respiratory system

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

contribute to the health effects of particulates. Particulates have been associated with increased respiratory diseases (asthma, bronchitis, and emphysema), cardiovascular disease (heart attack), and cancer.

Plant surfaces and growth rates may be adversely affected by particulate matter. Particulate air pollution also causes a wide range of damage to materials including corrosion of metals and electrical equipment and the soiling of textiles and buildings.

Sulfur Dioxide (SO₂)

Sulfur dioxide, (SO₂) is an atmospheric pollutant which results from combustion processes (mainly burning of fossil fuels containing sulfur compounds), refining of petroleum, manufacture of sulfuric acid, and smelting of ores containing sulfur. Reduction of sulfur dioxide pollution levels can generally be achieved through the use of low-sulfur content fuels or the use of chemical sulfur removal systems.

Once in the atmosphere, some sulfur dioxide can be oxidized (either photochemically or in the presence of a catalyst) to SO₃ (sulfur trioxide). In the presence of water vapor, SO₃ is readily converted to sulfuric acid (H₂SO₄) mist. Other basic oxides combine with SO₃ to form sulfate aerosols. Sulfuric acid droplets and other sulfates are thought to account for about 5 to 20 percent of the total suspended particulate matter in urban air. These compounds can be transported large distances and come back to earth as a major constituent of acid precipitation. Many of the resultant health problems attributed to SO₂ may be a result of the oxidation of SO₂ to other compounds.

The effects of SO₂ on health are irritation and inflammation of tissue that it directly contacts. Inhalation of SO₂ causes bronchial constriction resulting in an increased resistance to air flow, reduction of air volume, and an increase of respiratory rate and heart rate.

SO₂ can exacerbate pre-existing respiratory diseases (asthma, bronchitis, emphysema). The enhancement (synergism) by particulate

matter of the toxic response to SO₂ has been observed under conditions which would promote the conversion of SO₂ to H₂SO₄. The degree of enhancement is related to the concentration of particulate matter. A twofold to threefold increase of the irritant response to SO₂ is observed in the presence of particulate matter capable of oxidizing SO₂ to H₂SO₄.

H₂SO₄ inhalation causes an increase in the respiratory system's mucous secretions, which reduces the system's ability to remove particulates via mucociliary clearance. This can result in an increased incidence of respiratory infection.

Carbon Monoxide (CO)

The major source of carbon monoxide (CO) is motor vehicles. The USEPA has kept under its jurisdiction the regulation of emission control equipment on new motor vehicles while the State's responsibility for reducing excessive ambient carbon monoxide levels is exercised by developing transportation plans for congested urban areas.

The toxic effects of high concentrations of CO on the body are well known. Carbon monoxide is absorbed by the lungs and reacts with hemoglobin (the oxygen-carrying molecule in the blood) to form carboxyhemoglobin (COHb). This reaction reduces the oxygen-carrying capacity of blood because the affinity of hemoglobin for CO is over 200 times that for oxygen. The higher the percentage of hemoglobin bound up in the form of carboxyhemoglobin, the more serious is the health effect.

The level of COHb in the blood is directly related to the CO concentration of the inhaled air. For a given ambient air CO concentration, the COHb level in the blood will reach an equilibrium concentration after a sufficient time period. This equilibrium COHb level will be maintained in the blood as long as the ambient air CO level remains unchanged. However, the COHb level will slowly change in the same direction as the CO concentration of the ambient air as a new equilibrium of CO in the blood is established.

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

The lowest CO concentrations shown to produce adverse health effects result in aggravation of cardiovascular disease. Studies demonstrate that these concentrations have resulted in decreased exercise time before the onset of pain in the chest and extremities of individuals with heart or circulatory disease. Slightly higher CO levels have been associated with decreases in vigilance, the ability to discriminate time intervals, and exercise performance.

Evidence also exists indicating a possible relationship between CO and heart attacks, the development of cardiovascular disease, and irregular fetal development.

Studies on the existing ambient levels of CO do not indicate any adverse effects on vegetation, materials, or other aspects of human welfare.

Nitrogen Dioxide (NO₂)

Nitrogen gas (N₂) is an abundant and inert gas which makes up almost 80 percent of the Earth's atmosphere. In this form, it is harmless to humans and essential to plant metabolism. Due to its abundance in the air, it is a frequent reactant in many combustion processes. When combustion temperatures are extremely high, as in the burning of coal, oil, natural gas, and gasoline, atmospheric nitrogen gas may combine with molecular oxygen (O₂) to form various oxides of nitrogen (NO_x). Of these, nitric oxide (NO) and nitrogen dioxide (NO₂) are the most important contributors to air pollution; NO_x generally is used to represent these. Nitric oxide is a colorless and odorless gas. It is the primary form of NO_x resulting from the combustion process. NO_x contributes to haze and visibility reduction. NO_x is also known to cause deterioration and fading of certain fabrics and damage to vegetation. Depending on concentration and extent of exposure, plants may suffer leaf lesions and reduced crop yield.

Sensitivity of plants to NO_x depends on a variety of factors including species, time of day, light, stage of maturity, and the presence or absence of other air pollutants such as sulfur dioxide and ozone.

There is a lack of strong evidence associating health effects with most NO_x compounds. NO₂, a secondary derivative of atmospheric nitric oxide, however, has been clearly established as exerting detrimental effects on human health and welfare.

NO₂ can cause eye irritation at concentrations as low as 0.07 ppm. NO₂ can cause an increase in airway resistance, an increase in respiratory rate, an increase in sensitivity to bronchoconstrictors, a decrease in lung compliance, and an enhanced susceptibility to respiratory infections. NO₂ is a deep lung irritant capable of producing pulmonary edema if inhaled in sufficient concentrations. When NO₂ is inhaled in concentrations with other pollutants, the effects are additive.

NO_x may also react with water to form corrosive nitric acids, a major component of acid precipitation. Additionally, NO_x and various other pollutants (e.g., hydrocarbons) may react in the presence of sunlight to produce photochemical oxidants.

Lead (Pb)

Historically, atmospheric lead came primarily from combustion of leaded gasoline. However, the use of unleaded gas since 1975 has reduced mobile source lead emissions by over 90%. Currently stationary sources, such as lead smelters, battery manufacturers, and iron and steel producers can contribute significant amounts of lead to their immediate vicinity.

Lead is a stable compound which persists and accumulates both in the environment and in the human body. Lead enters the human body through ingestion and inhalation with consequent absorption into the blood stream and distribution to all body tissues. No safe level of lead in the blood has been identified. Clinical, epidemiological and toxicological studies have demonstrated exposure to lead has a broad range of health effects.

Since 1990, over 6,000 new health studies have been conducted. These studies have shown that children are the most susceptible to the

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

damaging effects of lead because they are more likely to ingest lead due to hand-to-mouth activity and early body development. Lead exposure has been found to interfere with the developing nervous system including the brain. This can potentially lead to intelligence quotient loss, poor academic achievement, permanent learning disabilities, and behavioral problems. These effects can persist into early adulthood.

Kidney and neurological cell damage has also been associated with lead exposure. Animal studies have demonstrated that lead can contribute to reduced fertility and birth defects.

Other potential effects from lead exposure are weakened immune systems, restlessness, headaches, increased blood pressure, and cardiovascular disease.

Illinois Ambient Air Quality Standards and Episode Levels

Consistent with the intent of the Environmental Protection Act of the State of Illinois, Illinois has adopted ambient air quality and episode standards that specify maximum permissible short-term and long-term concentrations of various contaminants in the atmosphere. Ambient air quality and episode standards are limits on atmospheric concentrations of air contaminants established for the purpose of protecting the public health and welfare.

The Illinois and National Ambient Air Quality Standards (NAAQS) consist of a primary and secondary standard for each pollutant (contaminant) as presented in **Table 1**. The Illinois Air Pollution Episode Levels are presented in **Table 2**. The primary standard and episode criterion represents the level of air quality which is necessary to protect the public health. Air entering the respiratory tract must not jeopardize health. Therefore, the air quality standards must, as a minimum, provide air which will not adversely affect, through acute or chronic symptoms, the public health.

The secondary standard defines the level of air quality which is necessary to protect the public welfare. This includes, among other things,

effects on crops, vegetation, wildlife, visibility, and climate, as well as effects on materials, economic values, and on personal comfort and well-being. The standards are legally enforceable limitations, and any person causing or contributing to a violation of the standards is subject to enforcement proceedings under the Environmental Protection Act. The standards have also been designed for use as a basis for the development of implementation plans by State and local agencies for the abatement and control of pollutant emissions from existing sources, and for the determination of air contaminant emission limitations to ensure that population, industry, and economic growth trends do not add to the region's air pollution problems.

Section 1: Air Pollutants – Sources, Health, and Welfare Effects

Table 1: Summary of National and Illinois Ambient Air Quality Standards				
Pollutant	Primary/ Secondary	Averaging Time	Level	Form
Carbon Monoxide	primary	8-hour	9 ppm	Not to be exceeded more than once per year
		1-hour	35 ppm	
Lead	primary and secondary	Rolling 3-month average	0.15 µg/m ³	Not to be exceeded
Nitrogen Dioxide	primary	1-hour	100 ppb	98th percentile, averaged over 3 years
	primary and secondary	Annual	53 ppb	Annual Mean
Ozone	primary and secondary	8-hour	0.070 ppm	Annual fourth-highest daily maximum 8-hr concentration, averaged over 3 years
Particle Pollution	PM _{2.5}	primary	Annual	12.0 µg/m ³
		secondary	Annual	15.0 µg/m ³
	PM ₁₀	primary and secondary	24-hour	35 µg/m ³
		24-hour	150 µg/m ³	98th percentile, averaged over 3 years
Sulfur Dioxide	primary	1-hour	75 ppb	99th percentile of 1-hour daily maximum concentrations, averaged over 3 years
	secondary	3-hour	0.5 ppm	Not to be exceeded more than once per year
PM _{2.5} standards are referenced to local conditions of temperature and pressure rather than standard conditions (760 mmHg and 25 degrees Celsius).				

Table 2: Illinois Air Pollution Episode Levels				
Pollutant	Advisory	Yellow Alert	Red Alert	Emergency
Particulate Matter (µg/m ³)	2-hour 420	24-hour 350	24-hour 420	24-hour 500
Sulfur Dioxide (ppm)	2-hour 0.30	4-hour 0.30	4-hour 0.35	4-hour 0.40
Carbon Monoxide (ppm)	2-hour 30	8-hour 15	8-hour 30	8-hour 40
Nitrogen Dioxide (ppm)	2-hour 0.40	1-hour 0.60 or 24-hour 0.15	1-hour 1.20 or 24-hour 0.30	1-hour 1.60 or 24-hour 0.40
Ozone (ppm)	1-hour 0.12	1-hour 0.20	1-hour 0.30	1-hour 0.50

Section 2: Statewide Summary of Air Quality

OZONE

Monitoring was conducted at 37 locations during the March-October "ozone season" and at least 75 percent data capture was obtained at all 37 sites.

Zion recorded the highest 1-hour concentration of 0.104 ppm. This compares with the highest concentration of 0.112 ppm in 2019 at Lisle. The highest value in the Metro-East area in 2020 was 0.095 ppm recorded at Wood River, compared with a high in 2019 of 0.108 ppm also at Wood River.

Data are also presented to compare with the current 8-hour standard as of 2016 of 0.070 ppm. The appropriate statistic for comparison with the 8-hour standard is the fourth highest value, which is averaged over a three-year period. There were 12 sites in Illinois that had a fourth-high value above 0.070 ppm in 2020 compared with two sites in 2019. The highest fourth-high value was 0.079 ppm at Northbrook and Cicero. The highest level in the Metro-East area was 0.069 ppm at Alton and Wood River. For the three-year period 2018-2020, 12 sites had a fourth-high average above 0.070 ppm (Table B4).

Figure 1 shows for each year the statewide average of each site's highest hourly ozone value for the ten-year period 2011-2020. The graph shows some year-to-year fluctuation with high years occurring during summers more favorable for ozone formation and low years in summers less conducive for ozone formation. The statewide average for 2020 was 0.085 ppm compared with 0.082 ppm in 2019 and 0.091 ppm in 2018.

Statewide, the total number of 1-hour excursion days in 2020 was zero compared with zero in 2019 and zero in 2018.





Figure 2 shows for each year the statewide annual average of the fourth highest 8-hour ozone value for the same period 2011-2020. The statewide average for 2020 was 0.069 ppm compared with 0.065 ppm in 2019 and 0.071 in 2018.

Section 2: Statewide Summary of Air Quality

PARTICULATE MATTER



Monitoring was conducted at 34 sites for PM_{2.5}. In 2020, no sites recorded an average above 12.0 ug/m³, the level of the annual standard. The statewide average of the annual averages was 8.7 ug/m³ in 2020 compared to 9.0 ug/m³ in 2019.

Figure 3 shows the trend of the statewide annual averages for PM_{2.5} for the period 2011-2020. There were nine exceedances of the 24-hour standard of 35 ug/m³ in 2020 compared with one exceedance in 2019 and two exceedances in 2018. The statewide peak of 62.9 ug/m³ was recorded in Joliet due to fireworks emissions. In 2020, the statewide 24-hour average was 19.9 ug/m³. This compares with 21.4 ug/m³ in 2019 and 21.3 ug/m³ in 2018.

In 2020 there were four sites monitoring PM₁₀. The statewide annual average was 30 ug/m³ compared with 27 ug/m³ in 2019 and 24 ug/m³ in 2018. The highest annual average was 37 ug/m³ in Lyons Township. The lowest annual was 20 ug/m³ at Northbrook.

For PM₁₀, the statewide average of the maximum 24-hour averages in 2020 was 98 ug/m³ compared with 77 ug/m³ in 2019 and 81 ug/m³ in 2018. **Figure 4** depicts this information for the period 2011-2020.

There was one exceedance of the 24-hour primary standard of 150 ug/m³. The highest 24-hour average was recorded in Lyons Township with a value of 159 ug/m³ compared with a high 24-hour value of 104 ug/m³ in Granite City in 2019.




Section 2: Statewide Summary of Air Quality

CARBON MONOXIDE

There were no exceedances of either the 1-hour primary standard of 35 ppm or the 8-hour primary standard of 9 ppm in 2020. The highest 1-hour average was 1.8 ppm recorded at the Lansing near-road location. The highest 8-hour average was 1.4 ppm also recorded at the Lansing near-road location.

Figure 5 shows the trend for the period 2011-2020 for the statewide average of the 1-hour and 8-hour high CO values. The statewide average of the 1-hour high was 1.0 ppm in 2020 compared with 1.8 ppm in 2019. The statewide average for the 8-hour high was 0.8 ppm in 2020 compared with 1.0 ppm in 2019.



There were no sites over the primary 1-hour standard of 75 ppb for the 2018-2020 period (Table B17).




Figure 6 shows the statewide trend for the maximum 24-hour averages for the period 2011-2020. The statewide average for 2020 was 6 ppb compared with the 2019 average of 6 ppb.

SULFUR DIOXIDE


There were no exceedances of the 1-hour primary standard of 75 ppb in 2020 compared with zero exceedances in 2019. There were no exceedances of the 3-hour secondary standard of 500 ppb in 2020. The highest 1-hour average was 61 ppb recorded in Decatur (Tate & Lyle) compared with 61 ppb in Mount Carmel in 2019. The statewide average of the 1-hour high in 2020 was 24 ppb. This compares with 26 ppb in 2019 and 34 ppb in 2018. The highest 3-hour average of 42 ppb was recorded in Decatur in 2020 compared with 45 ppb in Decatur in 2019.

Section 2: Statewide Summary of Air Quality

NITROGEN DIOXIDE

There were no violations of the annual primary standard of 53 ppb recorded in Illinois during 2020. The highest annual average of 16.5 ppb was recorded at the near-road site in Lansing. The statewide annual average for 2020 was 11.6 ppb compared with 12.5 ppb in 2019 and 12.2 ppb in 2018. There were no violations of the 1-hour primary standard, and there were also no violations in 2019. There were no sites over the 1-hour primary standard of 100 ppb for the 2018-2020 period compared to zero sites for the 2017-2019 period (Table B20).

Figure 7 depicts the trend of statewide annual averages from 2011-2020. There have been no violations of the annual standard since 1980.



LEAD

There were no violations of the rolling three-month maximum mean standard for the 2018 to 2020 period (Table B23).




Figure 8 shows the trend of the statewide maximum rolling three-month averages from 2011-2020. The decrease in 2013 was due to various controls having been implemented at facilities that have source-oriented monitors. The increase in 2019 was due to lead emission control problems at one facility in Granite City. The problems were addressed with the facility and corrective actions taken. Due to three-month averaging times, averages for January 2020 were affected as well (November 2019 – January 2020 average). All monitoring locations in the State have three-year maximum averages under the national standard for lead (Table B23). The statewide average for all sites was 0.057 ug/m³ in 2020 compared to 0.057 ug/m³ in 2019 and 0.027 ug/m³ in 2018.

Section 2: Statewide Summary of Air Quality

FILTER ANALYSIS RESULTS

The total suspended particulate samples were analyzed, in addition to lead, for specific metals. Several of the metals analyzed (arsenic, beryllium, cadmium, chromium, manganese, and nickel) have known toxic properties. Other metals such as iron can be used as tracers to help identify sources of high particulate values. There are currently no state or federal ambient air quality standards for these parameters.

The areas with the highest metals concentrations in Illinois are generally the heavily-industrialized areas of the Metro-East (Granite City and East St. Louis), south Chicago, and near source-oriented monitors. The highest 24-hour average for nickel was 0.041 ug/m³ measured in Granite City. The monitor at Granite City recorded the highest cadmium concentration with a 24-hour average of 0.007 ug/m³. The highest 24-hour chromium average was 0.245 ug/m³ recorded in Granite City. The highest iron and manganese values were also recorded in Granite City.

TOXIC COMPOUNDS

Sampling for toxic compounds other than metals (see Filter Analysis Section, **Table B24**) was conducted at Northbrook and Schiller Park. Most compounds were below the method detection limits. **Table B25** has a listing of various toxic compound maximums and annual averages.

Section 3: Air Quality Index

The Air Quality Index (AQI) is the national standard method for reporting air pollution levels to the public. An index such as the AQI is necessary because there are several air pollutants, each with different typical ambient concentrations and each with different levels of harm, and to report actual concentrations for all of them would be confusing. The AQI uses a single number and a short descriptor to define the air quality in an easy-to-remember and easy-to-understand way, taking all the pollutants into account.

The AQI is based on the short-term federal National Ambient Air Quality Standards (NAAQS), for six of the criteria pollutants, namely:

- Ozone (O_3)
- Sulfur dioxide (SO_2)
- Carbon monoxide (CO)
- Particulate matter (PM_{10})
- Particulate matter ($PM_{2.5}$)
- Nitrogen dioxide (NO_2)

In each case, the short-term primary NAAQS corresponds to 100 on the AQI scale – the top end of the Moderate category. The next concentration above the NAAQS would begin the Unhealthy for Sensitive Groups category at 101 on the AQI scale. **Table 3** lists all the AQI ranges and their descriptor categories. Each category corresponds to a different level of health concern. **Table 4** lists each AQI category and its corresponding meaning.

Unhealthy for Sensitive Groups occurs on occasion for 8-hour ozone, $PM_{2.5}$, and downwind of certain SO_2 sources. Unhealthy air quality is uncommon in Illinois, and Very Unhealthful air quality is rare. There has never been an occurrence of Hazardous air quality in Illinois.

The AQI is computed as follows: data from pollution monitors in an area are collected, and the AQI sub index for each pollutant is computed using formulas derived from the

index and concentration relations. Nomograms and tables are also available for this purpose. The data used are:

- O_3 estimate of the highest 8-hour average for that calendar day
- SO_2 the highest 1-hour or most recent 24-hour average
- CO the highest 8-hour average so far that calendar day
- PM_{10} the most recent 24-hour average
- $PM_{2.5}$ estimate of the 24-hour average for that calendar day
- NO_2 the highest 1-hour average

Continuous monitors are utilized for all the pollutants, including PM_{10} and $PM_{2.5}$.

Once all the sub-indices for the various pollutants have been computed, the highest is chosen.. That is the AQI for the area and the pollutant giving rise to it is the "critical pollutant." Thus if, for Anytown, Illinois, the following sub-indices were obtained:

O_3	= 45
SO_2	= 23
CO	= 19
PM_{10}	= 41
$PM_{2.5}$	= 61

Anytown's AQI for that day would be 61, which is in the Moderate category, and the critical pollutant would be particulates ($PM_{2.5}$). If data for one of the pollutants used in computing AQI is missing, the AQI is computed using the data available, ignoring the missing data. It occasionally happens that two pollutants have the same sub index; in such cases there are two critical pollutants.

The Illinois EPA issues an AQI forecast for 14 areas, or sectors, in Illinois (**Table 5**). These correspond to metropolitan areas with populations greater than 100,000.

Section 3: Air Quality Index

Table 3: Air Quality Index Categories

AQI Values	AQI Descriptor	Colors
<i>When the AQI is in this range:</i>	<i>...air quality conditions are:</i>	<i>...as symbolized by this color:</i>
0-50	Good	Green
51-100	Moderate	Yellow
101-150	Unhealthy for Sensitive Groups	Orange
151 to 200	Unhealthy	Red
201 to 300	Very Unhealthy	Purple
301 to 500	Hazardous	Maroon

Table 4: Air Quality Index Health Concerns

Air Quality Index Levels of Health Concern	Numerical Value	Meaning
Good	0 to 50	Air quality is considered satisfactory, and air pollution poses little or no risk.
Moderate	51 to 100	Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a very small number of people who are unusually sensitive to air pollution.
Unhealthy for Sensitive Groups	101 to 150	Members of sensitive groups may experience health effects. The general public is not likely to be affected.
Unhealthy	151 to 200	Everyone may begin to experience health effects; members of sensitive groups may experience more serious health effects.
Very Unhealthy	201 to 300	Health warnings of emergency conditions. The entire population is more likely to be affected.
Hazardous	301 to 500	Health alert: everyone may experience more serious health effects.

Section 3: Air Quality Index

Table 5: Air Quality Index Sectors in Illinois	
Sector	Coverage Area
Lake County	Lake County only
Chicago	All areas within the city limits of Chicago
North and West Suburbs	Parts of Cook, Du Page, and McHenry Counties north of I-290 (Eisenhower Expressway) and outside of the Chicago city limits
South and West Suburbs	Parts of Cook and Du Page Counties south of I-290 and outside of Chicago city limits
Will County/Joliet	Will County only
Aurora-Elgin	The eastern part of Kane County
Rockford	Approximately 10-mile diameter circle centered on downtown Rockford
Rock Island	The Illinois portion of the Quad Cities area
Peoria	Approximately 10-mile diameter circle centered on downtown Peoria in parts of Peoria, Woodford, and Tazewell Counties
Champaign	Champaign-Urbana Metropolitan Area
Normal	Bloomington-Normal Metropolitan Area
Decatur	Decatur Metropolitan Area
Springfield	Springfield Metropolitan Area
Metro-East St. Louis	The Illinois portion of the St. Louis Metropolitan Area. Approximately 15 miles wide east of the Mississippi River in Madison and St. Clair Counties

Section 3: Air Quality Index

Illinois EPA AQI forecasts and AQI information can be obtained on EPA's AirNow website at <http://www.airnow.gov>. The AirNow website shows estimated real-time AQI levels for all sectors in Illinois as well as other areas around the country. AQI information can further be obtained via e-mail and/or cell phones through the EnviroFlash program located at <http://illinois.enviroflash.info/signup.cfm>. The AirNow website and residents subscribed to EnviroFlash program can also receive alerts when high pollution levels are occurring or expected to occur. Additionally, Illinois AQI forecasts and current AQI levels are picked up and reported by various media outlets, weather websites, and electronic application programs.

2020 Illinois AQI Sector Summary

In order to present a more representative AQI, 24-hour calendar day FRM PM_{2.5} and PM₁₀ values from the total network were used to determine the percentages in **Figure 9** even though some of these values were not available for issuing the daily AQI.

Air quality was still in the "Good" and "Moderate" categories most often in 2020. All sectors had a higher frequency of "Good" than "Moderate", and all sectors had a higher frequency of "Moderate" than "Unhealthy for Sensitive Groups." Lake County, North & West Suburbs, Aurora-Elgin, Joliet/Will County, Rockford, Rock Island, Peoria, Champaign, Normal, Decatur, and Springfield sectors had 65 percent or more of the days in the "Good" category.

Within AQI sectors there were 72 occurrences of "Unhealthy for Sensitive Groups" air quality and 11 occurrences of "Unhealthy" air quality in 2020. The sector breakdown for "Unhealthy for Sensitive Groups" was 11 in Lake County, nine in Chicago, 17 in North & West Suburbs, 10 in South & West, seven in Aurora-Elgin, seven in Will County, one in Rockford, one in Rock Island, one in Peoria, three in Normal, two in Champaign and three in Metro-East. The sector breakdown for "Unhealthy" was one in Lake County, two in Chicago, one in North and West Suburbs, three in South & West


Suburbs, one in Aurora-Elgin and three in Joliet/Will County. **Figure 9** presents the AQI statistics for each sector. The pie chart shows the percent of days each sector was in a particular category.

In 2020, there were no ozone advisories issued in Illinois. An advisory is declared when ozone levels have reached the level of the former 1-hour standard (0.125 ppm) on a particular day. In the Chicago MSA there were 10 Air Pollution Action Days issued in 2020. This compares with zero in 2019.


Section 3: Air Quality Index

Figure 9: 2020 Air Quality Index Summaries by Sector


Chicago Sector - Lake County




Chicago Sector - Chicago




Chicago Sector - North & West Suburbs





Chicago Sector - South & West Suburbs



Aurora - Elgin




Joliet/Will County



Section 3: Air Quality Index


Figure 9: 2020 Air Quality Index Summaries by Sector




Section 3: Air Quality Index

Figure 9: 2020 Air Quality Index Summaries by Sector

Springfield



Metro-East (St. Louis)



Section 4: Statewide Summary of Point Source Emissions

Since the late 1970s, the Illinois EPA's Division of Air Pollution Control has maintained a database of stationary point source emissions for the entire State. 40 CFR 51.211 requires Illinois to include in its State Implementation Plan "... procedures for requiring owners or operators of stationary sources to maintain records of... a) Information on the nature and amount of emissions from the stationary source and b) other information as may be necessary..." The emission database maintained by the Division of Air Pollution Control has changed over time.

The current emissions inventory is known as the Integrated Comprehensive Environmental Management System (ICEMAN) and includes emission data on approximately 6,212 active sources (including 3,650 in the Registration of Smaller Sources, or ROSS, program) throughout the State. The ICEMAN data includes source addresses; source emission totals; permit data such as expiration date and status; emission unit data such as name, hours of operation, operating rate, fuel parameters, and emissions; control equipment data such as control device name, type, and removal efficiencies; and stack parameters. Reported emissions and Agency-calculated emissions are stored separately.

The group responsible for the entry of emission inventory data is the Regulatory Development Unit of the Air Quality Planning Section, and uses permit applications, the issued permit, and data reported on annual emissions reports to compile the inventory.

The following tables and graphs are an analysis of the emissions data contained in ICEMAN at the end of 2020. It is important to note emissions contained in the ICEMAN are not necessarily the actual emissions that entered the atmosphere. This is due to the fact that when an air pollution permit is applied for, the applicant provides maximum and average emission rates. The maximum emission rate reflects what the applicant believes the emission rate would be at maximum production. The average emission rate reflects emissions at the applicant's most

probable production rate. The Regulatory Development Unit has been updating its estimated emissions to more accurately reflect the reported emissions.

To calculate the distribution of emissions for the individual categories, the source classification code (SCC) field was used from the ICEMAN. The SCC is an eight-digit code that breaks emission units into logical categories. SCCs are provided by the USEPA.

To produce the following tables, the first three digits of the SCC were used. Only categories that contributed significantly to the overall total are listed in the following sections. The complete category breakdown can be found in Appendix C.

Section 4: Statewide Summary of Point Source Emissions

Volatile Organic Material

Figure 10
Volatile Organic Material
Emission Trend (1000s of Tons/Year)




Table 6: Volatile Organic Material Emissions - 2020

Category	Estimated Emissions (tons)	Category Contribution	Cumulative Percent
Food/Agriculture	9,718.2	25.61%	25.61%
Chemical Manufacturing	5,658.5	14.91%	40.52%
Surface Coating Operations	5,656.8	14.91%	55.43%
Petroleum Product Storage	2,368.0	6.24%	61.67%
Fuel Combustion	2,075.1	5.47%	67.14%
Printing/Publishing	1,947.2	5.13%	72.27%
Petroleum Industry	1,775.0	4.68%	76.94%
Rubber and Plastic Products	1,471.9	3.88%	80.82%
Mineral Products	939.8	2.48%	83.30%
Bulk Terminals/Plants	895.5	2.36%	85.66%
Organic Chemical Storage (large)	861.7	2.27%	87.93%
Secondary Metal Production	776.7	2.05%	89.98%
Fabricated Metal Products	710.6	1.87%	91.85%
Solid Waste Disposal	561.8	1.48%	93.33%
Organic Solvent Use	512.9	1.35%	94.68%
Petroleum Marketing/Transport	344.3	0.91%	95.59%
Organic Solvent Evaporation	341.4	0.90%	96.49%
All Other Categories	1,332.9	3.51%	100.00%

Section 4: Statewide Summary of Point Source Emissions

PM₁₀

Figure 11
PM₁₀ Emission Trend
(1000s of Tons/Year)




Table 7: Distribution of PM₁₀ Emissions – 2020

Category	Estimated Emissions (tons)	Category Contribution	Cumulative Percent
Food/Agriculture	5,756.1	29.51%	29.51%
Fuel Combustion	4,482.0	22.98%	52.49%
Mineral Products	3,597.1	18.44%	70.93%
Petroleum Industry	1,195.3	6.13%	77.05%
Chemical Manufacturing	992.5	5.09%	82.14%
Secondary Metal Production	851.6	4.37%	86.51%
Primary Metal Production	816.4	4.19%	90.69%
Solid Waste Disposal	508.3	2.61%	93.30%
Fabricated Metal Products	249.1	1.28%	94.58%
Surface Coating Operations	235.3	1.21%	95.78%
Process Cooling	230.5	1.18%	96.96%
All Other Categories	592.3	3.04%	100.00%

Section 4: Statewide Summary of Point Source Emissions

Carbon Monoxide

Figure 12
Carbon Monoxide Emission
Trend (1000s of Tons/Year)




Table 8: Distribution of Carbon Monoxide Emissions - 2020

Category	Estimated Emissions (tons)	Category Contribution	Cumulative Percent
Fuel Combustion	22,395.1	36.15%	36.15%
Primary Metal Production	21,676.6	34.99%	71.15%
In-Process fuel use	4,567.0	7.37%	78.52%
Mineral Products	3,039.8	4.91%	83.43%
Petroleum Industry	2,567.2	4.14%	87.57%
Solid Waste Disposal	2,074.7	3.35%	90.92%
Secondary Metal Production	1,893.8	3.06%	93.98%
Chemical Manufacturing	1,552.2	2.51%	96.48%
Food/Agriculture	1,201.8	1.94%	98.42%
Surface Coating Operations	237.0	0.38%	98.81%
Oil and Gas Production	220.5	0.36%	99.16%
All Other Categories	519.3	0.84%	100.00%

Section 4: Statewide Summary of Point Source Emissions

Sulfur Dioxide

Figure 13
Sulfur Dioxide Emission
Trend (1000s of Tons/Year)




Table 9: Distribution of Sulfur Dioxide Emissions - 2020

Category	Estimated Emissions (tons)	Category Contribution	Cumulative Percent
Fuel Combustion	61,085.5	82.76%	82.76%
Mineral Products	6,068.8	8.22%	90.98%
Petroleum Industry	1,629.0	2.21%	93.19%
Primary Metal Production	1,624.3	2.20%	95.39%
Food/Agriculture	1,301.3	1.76%	97.16%
Solid Waste Disposal	950.7	1.29%	98.44%
Chemical Manufacturing	850.4	1.15%	99.60%
All Other Categories	298.7	0.40%	100.00%

Section 4: Statewide Summary of Point Source Emissions

Nitrogen Oxides

Figure 14
Nitrogen Oxide Emission
Trend (1000s of Tons/Year)




Table 10: Distribution of Nitrogen Oxide Emissions – 2020

Category	Estimated Emissions (tons)	Category Contribution	Cumulative Percent
Fuel Combustion	41,843.9	71.79%	71.79%
Mineral Products	6,065.9	10.41%	82.19%
Petroleum Industry	3,571.4	6.13%	88.32%
Chemical Manufacturing	1,437.2	2.47%	90.79%
Primary Metal Production	1,269.9	2.18%	92.96%
Food/Agriculture	1,191.5	2.04%	95.01%
Solid Waste Disposal	740.3	1.27%	96.28%
Secondary Metal Production	691.6	1.19%	97.47%
Oil and Gas Production	555.8	0.95%	98.42%
Surface Coating Operations	468.3	0.80%	99.22%
All Other Categories	453.3	0.78%	100.00%

Appendix A: Air Sampling Network

Description of the Air Sampling Network

The Illinois air monitoring network is composed of instrumentation owned and operated by both the Illinois EPA and by cooperating local agencies. This network has been designed to measure ambient air quality levels throughout the State of Illinois following federal guidelines.

The network contains both continuous and non-continuous instruments. The continuous instruments operate throughout the year, while non-continuous instruments operate intermittently based on the schedule shown in **Table A1**. This is the official non-continuous sampling schedule used by the Illinois EPA during 2019.

The Illinois network is deployed along the lines described in the Illinois State Implementation Plan. An updated air monitoring plan is submitted to USEPA each year for review.

In accordance with USEPA air quality monitoring requirements as set forth in Title 40 of the Code of Federal Regulations, Part 58 (40 CFR 58), five types of monitoring stations are used to collect ambient air data. These include State and Local Air Monitoring Stations (SLAMS), National Air Monitoring Stations (NAMS), Photochemical Assessment Monitoring Stations (PAMS), Special Purpose Monitoring Stations (SPMS), and National Core Monitoring Stations (NCore). The types of stations are distinguished from one another on the basis of the general monitoring objectives they are designed to meet.

The SLAMS, NAMS, PAMS, SPMS, and NCore designations for the sites operated within the State of Illinois are provided in the Annual Network Plan, which can be found at epa.state.il.us/air/monitoring/index.html. All of the industrial sites are considered to be SPMS. **Table A2** is a summary of the distribution of pollutants through the years along with the total number of instruments and the total number of sites. The site directory is listed in **Table A3** and the monitoring directory is listed in **Table A4**.

Table A1
2020 Noncontinuous Sampling Schedule

JANUARY						
S	M	T	W	R	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

FEBRUARY						
S	M	T	W	R	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

MARCH						
S	M	T	W	R	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

APRIL						
S	M	T	W	R	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

MAY						
S	M	T	W	R	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

JUNE						
S	M	T	W	R	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

JULY						
S	M	T	W	R	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

AUGUST						
S	M	T	W	R	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

SEPTEMBER						
S	M	T	W	R	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			

OCTOBER						
S	M	T	W	R	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

NOVEMBER						
S	M	T	W	R	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

DECEMBER						
S	M	T	W	R	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

13 Every 6 Day Sampling Schedule **22** Every 3 Day Sampling Schedule

Appendix A: Air Sampling Network

1. **State/Local Air Monitoring Station (SLAMS) Network** - The SLAMS network is designed to meet a minimum of four basis monitoring objectives:
 - a. To determine the highest concentrations expected to occur in the area covered by the network.
 - b. To determine representative concentrations in areas of high population density.
 - c. To determine the air quality impact of significant sources or source categories.
 - d. To determine general background concentration levels.
2. **National Air Monitoring Station (NAMS) Network** - The NAMS network is a subset of stations selected from the SLAMS network with emphasis given to urban and multisource areas. The primary objectives of the NAMS network are:
 - a. To measure expected maximum concentrations.
 - b. To measure concentrations in areas where poor air quality is combined with high population exposure.
 - c. To provide data useable for the determination of national trends.
 - d. To provide data necessary to allow the development of nationwide control strategies.
3. **Photochemical Assessment Monitoring Station (PAMS) Network** - The PAMS network is required in serious, severe, and extreme ozone nonattainment areas to obtain detailed data for ozone, precursors (NO_x and VOC), and meteorology. NO_x and VOC sampling is required for the period June - August each year. Ozone sampling occurs during the ozone season, March - October. Network design is based on four monitoring types. In Illinois, PAMS are required in the Chicago metropolitan area only.
 - a. Type 1 sites are located upwind of the nonattainment area and are located to measure background levels of ozone and precursors coming into the area
 - b. Type 2 sites are located slightly downwind of the major source areas of ozone precursors.
 - c. Type 3 sites are located at the area of maximum ozone concentrations.
 - d. Type 4 sites are located at the domain edge of the nonattainment area and measure ozone and precursors leaving the area.
4. **Special Purpose Monitoring Station (SPMS) Network** - Any monitoring site that is not a designated SLAMS or NAMS is considered a special purpose monitoring station. Some of the SPMS network objectives are as follows:
 - a. To provide data as a supplement to stations used in developing local control strategies, including enforcement actions.

Appendix A: Air Sampling Network

- b. To verify the maintenance of ambient standards in areas not covered by the SLAMS/NAMS network.
 - c. To provide data on non-criteria pollutants.
5. **National Core Station (NCore) Network** - NCore is a multi-pollutant network that integrates several advanced measurement systems. In Illinois, Northbrook and Bondville are considered NCore sites. A few of the NCore network objectives are as follows:
- a. Support for development of emission strategies and accountability of emission strategy progress through tracking long-term trends of pollutants and their precursors.
 - b. Support of long-term health assessments that contribute to review of national standards.
 - c. Support to scientific studies ranging across technological, health, and atmospheric process disciplines.
 - d. Support to ecosystem assessments recognizing that national air quality networks benefit ecosystems assessments.

Appendix A: Air Sampling Network

Table A2
Distribution of Air Monitoring Equipment

Parameter	2020	2019	2018	2017	2016
Particulate Matter Federal Reference Method (PM _{2.5} FRM)	25	25	24	27	27
PM _{2.5} Federal Equivalent Method (PM _{2.5} FEM)	17	17	16	8	8
PM _{10-2.5} (PM Coarse)	1	1	1	0	0
PM _{2.5} Air Quality Index (non-FEM)	7	7	7	9	9
PM _{2.5} Speciation	4	4	4	4	5
Particulate Matter (PM ₁₀)	5	5	5	5	5
Lead (Pb)	5	5	5	7	7
Sulfur Dioxide (SO ₂)	14	14	14	10	13
Nitrogen Dioxide (NO ₂)	7	7	5	5	6
Total Reactive Nitrogen (NO _y)	2	2	2	2	2
Ozone (O ₃)	37	37	37	37	37
Carbon Monoxide (CO)	3	4	3	3	3
Volatile Organic Compounds	2	2	2	2	2
Semi Volatile Organic Compounds	1	1	1	1	1
Semi Non-Methane Organic Compounds	1	1	1	1	1
Carbonyls	2	2	2	2	2
Meteorology	11	11	17	19	20
Total Instruments	144	145	146	142	148
Total Sites	64	64	63	64	64

Appendix A: Air Sampling Network

Statewide Air Monitoring Site Locations - 2020

See the 2020 Site Directory (Table A3) for additional information.




Table A3

Site Directory

Site Map ID	AQS ID	County	City	Address	Latitude Longitude	Owner / Operator
1	17-001-0007	Adams	Quincy	John Wood Comm. College 1301 South 48th St.	+39.91540937 -91.33586832	IL EPA
2	17-019-1001	Champaign	Bondville	State Water Survey Township Rd. 500 E.	+40.052780 -88.372510	IL EPA/US EPA
3	17-019-0006	Champaign	Champaign	Ameren Substation 904 N. Walnut	+40.1237962 -88.229531	IL EPA
4	17-019-0007	Champaign	Thomasboro	North Thomas St.	+40.244913 -88.188519	IL EPA
5	17-023-0001	Clark	West Union	416 S. State Highway 1 & West Union	+39.210883 -87.668416	Indiana DEP
6	17-031-0001	Cook	Alsip	Village Garage 4500 W. 123rd St.	+41.6709919 -87.7324569	CCDES
7	17-031-0076	Cook	Chicago	Com Ed Maintenance Bldg. 7801 Lawndale	+41.75139998 -87.71348815	CCDES
8	17-031-0219	Cook	Chicago	Kennedy Near-road #2 Kennedy Expy. & W. Webster Ave.	+41.920681 -87.674425	IL EPA
9	17-031-0052	Cook	Chicago	Mayfair Pump Station 4850 Wilson Ave.	+41.96548483 -87.74992806	CCDES
10	17-031-0110	Cook	Chicago	Perez Elementary School 1241 19th St.	+41.855771 -87.657932	CCDES
11	17-031-0032	Cook	Chicago	South Water Filtration Plant 3300 E. Cheltenham Pl.	+41.75583241 -87.54534967	CCDES
12	17-031-0057	Cook	Chicago	Springfield Pump Station 1745 N. Springfield Ave.	+41.912739 -87.722673	CCDES
13	17-031-1003	Cook	Chicago	Taft High School 6545 W. Hurlbut St	+41.98433233 -87.7920017	CCDES
14	17-031-0022	Cook	Chicago	Washington High School 3535 E. 114th St.	+41.68716544 -87.53931548	CCDES
15	17-031-4002	Cook	Cicero	Cook County Trailer 1820 S. 51st Ave	+41.85524313 -87.7524697	CCDES
16	17-031-6005	Cook	Cicero	Liberty School 13th St. & 50th Ave.	+41.86442642 -87.74890238	CCDES
17	17-031-4007	Cook	Des Plaines	Regional Office Building 9511 W. Harrison St	+42.06028469 -87.86322543	IL EPA
18	17-031-7002	Cook	Evanston	Water Pumping Station 531 E. Lincoln	+42.062053 -87.675254	IL EPA
19	17-031-0119	Cook	Lansing	Kingery Near-road #1 Kingery Expy. & Torrence Ave.	+41.578603 -87.557392	IL EPA
20	17-031-1601	Cook	Lemont	Cook County Trailer 729 Houston	+41.66812034 -87.99056969	CCDES
21	17-031-1016	Cook	Lyons Township	Village Hall 50th St & Glencoe	+41.801180 -87.832349	IL EPA
22	17-031-4201	Cook	Northbrook	Northbrook Water Plant 750 Dundee Rd.	+42.13999619 -87.79922692	IL EPA
23	17-031-3103	Cook	Schiller Park	IEPA Trailer 4743 Mannheim Rd.	+41.96519348 -87.87626473	IL EPA
24	17-031-3301	Cook	Summit	Graves Elementary School 60th St. & 74th Ave.	+41.78276601 -87.80537679	CCDES

Table A3
Site Directory

Site Map ID	AQS ID	County	City	Address	Latitude Longitude	Owner / Operator
25	17-043-6001	DuPage	Lisle	Morton Arboretum Route 53	+41.81304939 -88.0728269	IL EPA
26	17-043-4002	DuPage	Naperville	City Hall 400 S. Eagle St.	+41.77107094 -88.15253365	IL EPA
27	17-049-1001	Effingham	Effingham	Central Grade School 10421 N. US Hwy. 45	+39.06715932 -88.54893401	IL EPA
28	17-065-0002	Hamilton	Knight Prairie	Ten Mile Creek DNR Office State Route 14	+38.08215516 -88.6249434	IL EPA
29	17-083-0117	Jerseyville	Jerseyville	21965 Maple Summit Rd.	+39.101439 -90.344494	IL EPA
30	17-085-9991	Jo Daviess	Stockton	10952 E. Parker Rd.	+42.2869 -89.9997	US EPA
31	17-089-0007	Kane	Aurora	Health Department 1240 N. Highland	+41.78471651 -88.32937361	IL EPA
32	17-089-0005	Kane	Elgin	Larsen Junior High School 665 Dundee Rd.	+42.04914776 -88.27302929	IL EPA
33	17-089-0003	Kane	Elgin	McKinley School 258 Lovell St.	+42.050403 -88.28001471	IL EPA
34	17-097-1007	Lake	Zion	Camp Logan Illinois Beach State Park	+42.4675733 -87.81004705	IL EPA
35	17-099-0007	La Salle	Oglesby	308 Portland Ave.	+41.29301454 -89.04942498	IL EPA
36	17-115-0013	Macon	Decatur	IEPA Trailer 2200 N. 22nd	+39.866933 -88.925452	IL EPA
37	17-115-0117	Macon	Decatur	ADM 2550 N. Brush College Rd.	+39.880404 -88.894488	ERM Inc.
38	17-115-0217	Macon	Decatur	Tate & Lyle North 899 N. Folk St.	+39.850712 -88.933635	ERM Inc.
39	17-115-0317	Macon	Decatur	Tate & Lyle South 2200 E. El Dorado St.	+39.846856 -88.923323	ERM Inc.
40	17-117-0002	Macoupin	Nilwood	IEPA Trailer Heaton & Dubois	+39.39607533 -89.80973892	IL EPA
41	17-119-0120	Madison	Alton	Horace Mann School 2708 Edwards St.	+38.901316 -90.146211	IL EPA
42	17-119-2009	Madison	Alton	SIU Dental Clinic 1700 Annex St.	+38.90308534 -90.14316803	IL EPA
43	17-119-0010	Madison	Granite City	Air Products 15th & Madison	+38.69443831 -90.15395426	IL EPA
44	17-119-1007	Madison	Granite City	Fire Station #1 23rd & Madison	+38.70453426 -90.13967484	IL EPA
45	17-119-0024	Madison	Granite City	Gateway Medical Center 2100 Madison Ave.	+38.7006315 -90.14476267	IL EPA
46	17-119-9991	Madison	Highland	5403 State Rd. 160	+38.8690 -89.6228	US EPA
47	17-119-1009	Madison	Maryville	200 West Division	+38.72657262 -89.95996251	IL EPA
48	17-119-3007	Madison	Wood River	Water Treatment Plant 54 N. Walcott	+38.86066947 -90.10585111	IL EPA
49	17-111-0001	McHenry	Cary	Cary Grove High School 1st St. & Three Oaks Rd.	+42.22144166 -88.24220734	IL EPA

Table A3
Site Directory

Site Map ID	AQS ID	County	City	Address	Latitude Longitude	Owner / Operator
50	17-113-2003	McLean	Normal	ISU Physical Plant Main & Gregory	+40.51873537 -88.99689571	IL EPA
51	17-143-0037	Peoria	Peoria	City Office Building 613 N.E. Jefferson	+40.697326 -89.584084	IL EPA
52	17-143-0024	Peoria	Peoria	Fire Station #8 MacArthur & Hurlburt	+40.68742038 -89.60694277	IL EPA
53	17-143-1001	Peoria	Peoria Heights	Peoria Heights High School 508 E. Glen Ave.	+40.74550393 -89.58586902	IL EPA
54	17-157-0001	Randolph	Houston	IEPA Trailer Hickory Grove & Fallview	+38.17627761 -89.78845862	IL EPA
55	17-161-3002	Rock Island	Rock Island	Rock Island Arsenal 32 Rodman Ave.	+41.51472697 -90.51735026	IL EPA
56	17-167-0012	Sangamon	Springfield	Agricultural Building State Fair Grounds	+39.83192087 -89.64416359	IL EPA
57	17-167-0014	Sangamon	Springfield	Illinois Building State Fair Grounds	+39.831522 -89.640926	IL EPA
58	17-163-0010	St. Clair	East St. Louis	RAPS Trailer 13th & Tudor	+38.61203448 -90.16047663	IL EPA
59	17-179-0004	Tazewell	Pekin	Fire Station #3 272 Derby	+40.55643203 -89.65402083	IL EPA
60	17-185-0001	Wabash	Mount Carmel	Division St.	+38.397276 -87.773631	Indiana DEP
61	17-197-1011	Will	Braidwood	Com Ed Training Center 36400 S. Essex Rd.	+41.22153707 -88.19096718	IL EPA
62	17-197-1002	Will	Joliet	Pershing Elementary School Midland & Campbell Sts.	+41.52688509 -88.11647381	IL EPA
63	17-201-2001	Winnebago	Loves Park	Maple Elementary School 1405 Maple Ave.	+42.33498222 -89.0377748	IL EPA
64	17-201-0118	Winnebago	Rockford	Fire Department 204 S. 1 st St.	+42.2670002 -89.089170	IL EPA

Table A4
Monitoring Directory

AQS ID	City	CO	NOy	NO2	Ozone	PM10	PM Coarse	PM2.5 FRM	PM2.5 FEM	PM2.5 AQI	PM2.5 Speciation	SO2	VOC	Toxics	TSP Pb, Metals	Meteorological
17-001-0007	Quincy															
17-019-0006	Champaign N. Walnut															
17-019-0007	Thomasboro															
17-019-1001	Bondville	T										T				
17-023-0001	West Union															
17-031-0001	Alsip															
17-031-0022	Chicago Washington High School					C										
17-031-0032	Chicago South Water Filtration															
17-031-0052	Chicago Mayfair Pump Station															
17-031-0057	Chicago Springfield Pump Station															
17-031-0076	Chicago Com Ed Maintenance															
17-031-0110	Chicago Perez Elementary															
17-031-0119	Lansing Kingery near-road #1															
17-031-0219	Chicago Kennedy near-road #2															
17-031-1003	Chicago Taft High School															
17-031-1016	Lyons Township					C										
17-031-1601	Lemont															
17-031-3103	Schiller Park															
17-031-3301	Summit															
17-031-4002	Cicero Cook County Trailer															
Active Monitor	Site/Monitor Installed	Site/Monitor Removed			C = Continuous PM ₁₀ , T = Trace level											

Table A4
Monitoring Directory

AQS ID	City	CO	NOy	NO2	Ozone	PM10	PM Coarse	PM2.5 FRM	PM2.5 FEM	PM2.5 AQI	PM2.5 Speciation	SO2	VOC	Toxics	TSP Pb, Metals	Meteorological
17-031-4007	Des Plaines															
17-031-4201	Northbrook	T										T				
17-031-6005	Cicero Liberty School															
17-031-7002	Evanston															
17-043-4002	Naperville															
17-043-6001	Lisle															
17-049-1001	Effingham															
17-065-0002	Knight Prairie															
17-083-0117	Jerseyville															
17-085-9991	Stockton															
17-089-0003	Elgin McKinley School															
17-089-0005	Elgin Larsen Jr. High School															
17-089-0007	Aurora															
17-097-1007	Zion															
17-099-0007	Oglesby															
17-111-0001	Cary															
17-113-2003	Normal															
17-115-0013	Decatur IEPA Trailer															
17-115-0117	Decatur ADM															
17-115-0217	Decatur Tate & Lyle North															
Active Monitor	Site/Monitor Installed	Site/Monitor Removed			T = Trace level											

Table A4
Monitoring Directory

AQS ID	City	CO	NOy	NO2	Ozone	PM10	PM Coarse	PM2.5 FRM	PM2.5 FEM	PM2.5 AQI	PM2.5 Speciation	SO2	VOC	Toxics	TSP Pb, Metals	Meteorological
17-115-0317	Decatur Tate & Lyle South															
17-117-0002	Nilwood															
17-119-0008	Alton Clara Barton Elementary															
17-119-0120	Alton Horace Mann School							*								
17-119-2009	Alton SIU Dental Clinic							*								
17-119-0010	Granite City Air Products															
17-119-0024	Granite City Gateway Medical Center															
17-119-1007	Granite City Fire Station #1															
17-119-1009	Maryville															
17-119-3007	Wood River															
17-119-9991	Highland															
17-143-0024	Peoria Fire Station #8															
17-143-0037	Peoria City Office Building															
17-143-1001	Peoria Heights															
17-157-0001	Houston															
17-161-3002	Rock Island															
17-163-0010	East St. Louis															
17-167-0012	Springfield Agricultural Building															
17-167-0014	Springfield Illinois Building															
17-179-0004	Pekin															
Active Monitor	Site/Monitor Installed	Site/Monitor Removed			*Alton PM2.5 to be moved from SIU Dental Clinic to Horace Mann School in 2021.											

Table A4
Monitoring Directory

AQS ID	City	CO	NOy	NO2	Ozone	PM10	PM Coarse	PM2.5 FRM	PM2.5 FEM	PM2.5 AQI	PM2.5 Speciation	SO2	VOC	Toxics	TSP Pb, Metals	Meteorological
17-185-0001	Mount Carmel															
17-197-1002	Joliet Pershing Elementary															
17-197-1011	Braidwood															
17-201-0118	Rockford Fire Department															
17-201-2001	Loves Park															
Active Monitor	Site/Monitor Installed	Site/Monitor Removed														

Air Quality Data Interpretation

In order to provide a uniform procedure for determining whether a sufficient amount of air quality data has been collected by a sensor in a given time period (year, quarter, month, day, etc.) to accurately represent air quality during that time period, a minimum statistical selection criteria was developed.

In order to calculate an annual average for non-continuous parameters, a minimum of 75% of the data that was scheduled to be collected must be available, i.e., 45 samples per year for an every-six-day schedule (total possible of 60 or 61 samples). Additionally, in order to have proper quarterly balance, each site on an every sixth day schedule should have at least 10 samples per calendar quarter. This provides for a 20% balance in each quarter if the minimum required annual sampling is achieved.

PM₁₀ and PM_{2.5} samplers operate on one of three sampling frequencies:

- Every-day sampling (68 samples required each quarter for 75% data capture)
- Every-third-day sampling (23 samples required each quarter for 75% data capture)
- Every-six-day sampling (12 samples required each quarter for 75% data capture).

To calculate an annual PM₁₀ or PM_{2.5} mean, arithmetic means are calculated for each quarter in which valid data is recorded in at least 75% of the possible sampling periods. The annual mean is then the arithmetic average of the four quarterly means.

To determine an annual average for continuous data 75% of the total possible yearly observations are necessary, i.e., a minimum of 6570 hours (75% of the hours available) are needed. In order to provide a balance between the respective quarters, each quarter should have at least 1300 hours which is 20% of the 75% minimum annual requirement. To calculate

quarterly averages at sites which do not meet the annual criteria, 75% of the total possible observations in a quarter are needed, i.e., a minimum of 1647 hours of 2200 hours available. Monthly averages also require 75% of the total possible observations in a month, i.e., 540 hours as a minimum. Additionally, for short-term running averages (24-hour, 8-hour, and 3-hour) 75% of the data during the particular time period is needed, i.e., 18 hours for a 24-hour average, six hours for an 8-hour average and three hours for a 3-hour average.

For ozone, a valid 8-hour average has at least six valid 1-hour averages within the 8-hour period. The daily maximum 8-hour ozone concentration is based on 17 consecutive moving 8-hour periods in each day, beginning with the 8-hour period from 7:00 a.m. to 3:00 p.m. and ending with the 8-hour period from 11:00 p.m. to 7:00 a.m. The daily maximum value is considered valid if 8-hour averages are available for at least 13 of the 17 consecutive moving 8-hour periods, or if the daily maximum value is greater than the level of the NAAQS. Complete sampling over a three-year period requires an average of 90% valid days with each year having at least 75% valid days.

Data listed as not meeting the minimum statistical selection criteria in this report were so noted after evaluation using the criteria above. Although short term averages (3, 8, 24 hours) have been computed for certain sites not meeting the annual criteria, these averages may not be representative of an entire year's air quality. In certain circumstances where even the 75% criteria is met, the number and/or magnitude of short-term averages may not be directly comparable from one year to the next because of seasonal distributional differences.

For summary purposes, the data is expressed in the number of figures to which the raw data is validated. Extra figures may be carried in the averaging technique, but the result is rounded to the appropriate number of figures. For example, the values 9, 9, and 10 are


Appendix B: Air Quality Data Summary Tables

averaged to give 9; whereas the values 9.0, 9.0, and 10.0 are averaged to 9.3. The raw data itself should not be expressed to more significant figures than the sensitivity of the monitoring methodology allows.

In comparing data to the various air quality standards, the data are implicitly rounded to the number of significant figures specified by that standard. For example, to exceed the 0.15 $\mu\text{g}/\text{m}^3$ three-month lead standard, a three-month average value must be 0.155 $\mu\text{g}/\text{m}^3$ or higher; to exceed the 9 ppm CO 8-hour standard, an 8-hour average must be 9.5 ppm or higher. Peak averages, though, will be expressed to the number of significant figures appropriate to that monitoring methodology.

The NAAQS for CO has a short-term standard for ambient air concentrations not to be exceeded more than once per year. SO_2 has a 1-hour standard which is the three-year average of each year's 99th percentile values. NO_2 has a 1-hour standard which is the three-year average of each year's 98th percentile values. PM_{10} has a 24-hour standard which cannot average more than one exceedance over a three-year period (in three years). $\text{PM}_{2.5}$ has a 24-hour standard which is a three-year average of each year's 98th percentile values. In the case of ozone, the 8-hour standard is concentration-based and as such is the average of the fourth highest value each year over a three-year period. The standards are promulgated in this manner in order to protect the public from excessive levels of pollution both in terms of acute and chronic health effects.

The following data tables detail and summarize air quality in Illinois. The tables of short-term exceedances list those sites which exceeded any of the short-term primary standards (24 hours or less). The detailed data tables list averages and peak concentrations for all monitoring sites in Illinois.



	Site ID	Site Name
1.	170010007	Quincy
2.	170190007	Thomasboro
3.	170191001	Bondville
4.	170230001	West Union
5.	170310001	Alsip
6.	170310032	Chicago – South Water Filtration
7.	170313103	Schiller Park
8.	170311601	Lemont
9.	170311003	Chicago – Taft High School
10.	170310076	Chicago – Com Ed Maint. Bldg.
11.	170314002	Cicero
12.	170314007	Des Plaines
13.	170314201	Northbrook
14.	170317002	Evanston
15.	170436001	Lisle
16.	170491001	Effingham
17.	170650002	Knight Prairie
18.	170831001	Jerseyville
19.	170859991	Stockton
20.	170890005	Elgin
21.	170971007	Zion
22.	171110001	Cary
23.	171132003	Normal
24.	171150013	Decatur
25.	171170002	Nilwood
26.	171190008	Alton
27.	171191009	Maryville
28.	171193007	Wood River
29.	171199991	Highland
30.	171430024	Peoria
31.	171431001	Peoria Heights
32.	171570001	Houston
33.	171613002	Rock Island
34.	171630010	East St. Louis
35.	171670014	Springfield
36.	171971011	Braidwood
37.	172012001	Loves Park

Total Over 0.12 ppm	0	
Total Days Over 0.12 ppm	0	

Table B2
8-Hour Ozone Exceedances

EXCEEDANCES OF THE 8-HOUR PRIMARY STANDARD OF 0.070 PPM						
Date	City	Concentration		Date	City	Concentration
6/4	Alsip	0.072		6/19	Cicero	0.086
	Chicago-SWFP	0.074			Chicago-Com Ed	0.084
	Evanston	0.072			Evanston	0.084
	Lemont	0.071			Des Plaines	0.083
	Northbrook	0.071			Northbrook	0.082
6/5	Chicago-SWFP	0.087			Chicago-Taft	0.077
	Evanston	0.083			Alsip	0.076
	Northbrook	0.082			Lemont	0.074
	Zion	0.073		6/25	Wood River	0.074
	Des Plaines	0.072			Alton	0.072
6/8	Cary	0.082		6/27	Evanston	0.074
	Elgin	0.073			Zion	0.071
	Des Plaines	0.071		7/1	Elgin	0.077
6/16	Cary	0.076			Alsip	0.073
	Elgin	0.073			Lisle	0.073
6/17	Elgin	0.081			Lemont	0.071
	Cary	0.079		7/2	Lemont	0.072
	Lemont	0.078		7/3	Lemont	0.082
	Alsip	0.077			Northbrook	0.078
	Chicago-SWFP	0.077			Lisle	0.077
	Lisle	0.077			Alsip	0.076
	Zion	0.074			Chicago-SWFP	0.075
	Evanston	0.073			Cicero	0.074
	Northbrook	0.073			Cary	0.073
	Des Plaines	0.072			Elgin	0.073
6/18	Elgin	0.090			Zion	0.073
	Lemont	0.088			Evanston	0.072
	Lisle	0.087			Normal	0.072
	Chicago-SWFP	0.085			Braidwood	0.070
	Cary	0.084		7/4	Normal	0.081
	Des Plaines	0.084			Thomasboro	0.078
	Alsip	0.083			Peoria Heights	0.075
	Chicago-Taft	0.081			Bondville	0.072
	Cicero	0.080		7/5	Normal	0.073
	Evanston	0.080		7/6	Des Plaines	0.088
	Northbrook	0.080			Lemont	0.086
	Zion	0.078			Cicero	0.082
	Jerseyville	0.074			Northbrook	0.079
	Braidwood	0.073			Chicago-Com Ed	0.076
	Thomasboro	0.073			Zion	0.076
	Schiller Park	0.072			Alsip	0.073
	Loves Park	0.071			Cary	0.073
6/19	Zion	0.099			Schiller Park	0.072
	Chicago-SWFP	0.091			Evanston	0.071

Total Over 0.070 ppm	113
Total Days Over 0.070 ppm	24

Table B3
Ozone Highs

AQS ID	City	Number Of Days 8-Hour Greater Than 0.070 ppm			Fourth Highest Samples 1-Hour (ppm)				Fourth Highest Samples 8-Hour (ppm)			
		2020	2019	2018								
17-001-0007	Quincy	0	0	0	0.080	0.070	0.068	0.068	0.065	0.064	0.064	0.064
17-019-0007	Thomasboro	2	0	4	0.082	0.075	0.075	0.073	0.078	0.073	0.070	0.069
17-019-1001	Bondville	1	0	1	0.081	0.071	0.068	0.064	0.072	0.068	0.062	0.062
17-023-0001	West Union	0	0	1	0.070	0.066	0.064	0.063	0.069	0.062	0.061	0.060
17-031-0001	Alsip	10	3	10	0.091	0.087	0.084	0.083	0.083	0.077	0.076	0.076
17-031-0032	Chicago South Water Filtration	6	4	7	0.102	0.094	0.094	0.087	0.091	0.087	0.085	0.077
17-031-0076	Chicago Com Ed Maintenance	2	1	8	0.093	0.088	0.082	0.078	0.084	0.076	0.069	0.068
17-031-1003	Chicago Taft High School	6	2	6	0.097	0.089	0.085	0.083	0.081	0.081	0.078	0.077
17-031-1601	Lemont	8	3	2	0.099	0.095	0.094	0.090	0.088	0.086	0.082	0.078
17-031-3103	Schiller Park	2	1	1	0.086	0.081	0.078	0.076	0.072	0.072	0.069	0.068
17-031-4002	Cicero Cook County Trailer	6	0	5	0.096	0.093	0.090	0.089	0.086	0.082	0.080	0.079
17-031-4007	Des Plaines	6	1	10	0.101	0.097	0.090	0.083	0.088	0.084	0.083	0.072
17-031-4201	Northbrook	11	0	10	0.093	0.090	0.088	0.087	0.082	0.082	0.080	0.079
17-031-7002	Evanston	10	3	12	0.092	0.090	0.085	0.084	0.084	0.083	0.080	0.074
17-043-6001	Lisle	4	3	6	0.093	0.085	0.084	0.081	0.087	0.077	0.077	0.073
17-049-1001	Effingham	0	0	1	0.073	0.070	0.068	0.066	0.065	0.064	0.064	0.062
17-065-0002	Knight Prairie	0	1	3	0.073	0.071	0.071	0.070	0.070	0.067	0.067	0.067
17-083-1001	Jerseyville	1	2	3	0.083	0.075	0.073	0.071	0.074	0.066	0.065	0.062
17-085-9991	Stockton	0	0	2	0.072	0.071	0.069	0.067	0.066	0.064	0.064	0.063
17-089-0005	Elgin Larsen Jr. High School	7	4	5	0.087	0.086	0.083	0.082	0.090	0.081	0.077	0.073
17-097-1007	Zion	11	2	8	0.104	0.092	0.087	0.086	0.099	0.078	0.078	0.076
17-111-0001	Cary	8	2	8	0.097	0.089	0.087	0.084	0.084	0.082	0.079	0.076
17-113-2003	Normal	3	0	1	0.083	0.076	0.076	0.075	0.081	0.073	0.072	0.070
17-115-0013	Decatur IEPA Trailer	0	0	3	0.071	0.070	0.069	0.068	0.067	0.067	0.066	0.065
17-117-0002	Nilwood	0	0	3	0.072	0.071	0.070	0.067	0.068	0.066	0.065	0.063

Table B3
Ozone Highs

AQS ID	City	Number Of Days 8-Hour Greater Than 0.070 ppm			Fourth Highest Samples 1-Hour (ppm)				Fourth Highest Samples 8-Hour (ppm)			
		2020	2019	2018								
17-119-0120	Alton	1	3	5	0.087	0.085	0.081	0.075	0.072	0.070	0.069	0.069
17-119-1009	Maryville	1	0	6	0.087	0.081	0.075	0.075	0.072	0.069	0.068	0.067
17-119-3007	Wood River	3	3	4	0.095	0.090	0.084	0.083	0.074	0.072	0.072	0.069
17-119-9991	Highland	0	0	4	0.080	0.073	0.072	0.071	0.068	0.067	0.067	0.066
17-143-0024	Peoria Fire Station #8	0	1	2	0.071	0.071	0.070	0.067	0.068	0.065	0.065	0.064
17-143-1001	Peoria Heights	1	1	3	0.078	0.077	0.075	0.073	0.075	0.070	0.070	0.070
17-157-0001	Houston	0	0	1	0.073	0.070	0.068	0.066	0.070	0.064	0.061	0.061
17-161-3002	Rock Island	0	2	1	0.075	0.070	0.068	0.067	0.069	0.065	0.065	0.063
17-163-0010	East St. Louis	0	1	5	0.078	0.077	0.072	0.072	0.070	0.068	0.065	0.065
17-167-0014	Springfield	0	0	1	0.072	0.071	0.071	0.071	0.069	0.068	0.068	0.067
17-197-1011	Braidwood	2	0	4	0.086	0.075	0.075	0.073	0.079	0.073	0.069	0.067
17-201-2001	Loves Park	1	0	3	0.077	0.075	0.073	0.072	0.071	0.069	0.068	0.067
Statewide Average					0.085	0.080	0.077	0.075	0.077	0.072	0.071	0.069
Total Over 0.070 ppm		113	43	159								
Total Days Over 0.070 ppm		24	16	26								

Table B4
Ozone Design Values



AQS ID	City	Fourth High 8-Hour Concentrations (ppm)					Design Values* (ppm)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-001-0007	Quincy	0.064	0.062	0.063	0.065	0.061	0.063	0.063	0.063
17-019-0007	Thomasboro	0.069	0.062	0.072	0.067	0.066	0.067	0.067	0.068
17-019-1001	Bondville	0.062	0.058	0.064	0.067	0.066	0.061	0.063	0.065
17-023-0001	West Union	0.060	0.060	0.066	0.067	0.066	0.062	0.064	0.066
17-031-0001	Alsip	0.076	0.070	0.079	0.078	0.075	0.075	0.075	0.077
17-031-0032	Chicago South Water Filtration	0.077	0.071	0.076	0.074	0.077	0.074	0.073	0.075
17-031-0076	Chicago Com Ed Maintenance	0.068	0.065	0.074	0.078	0.075	0.069	0.072	0.075
17-031-1003	Chicago Taft High School	0.077	0.069	0.073	0.060	0.075	0.073	0.067	0.069
17-031-1601	Lemont	0.078	0.068	0.068	0.070	0.073	0.071	0.068	0.070
17-031-3103	Schiller Park	0.068	0.064	0.065	0.061	0.067	0.065	0.063	0.064
17-031-4002	Cicero Cook County Trailer	0.079	0.064	0.072	0.068	0.076	0.071	0.068	0.072
17-031-4007	Des Plaines	0.072	0.066	0.075	0.071	0.076	0.071	0.070	0.074
17-031-4201	Northbrook	0.079	0.069	0.083	0.070	0.079	0.077	0.074	0.077
17-031-7002	Evanston	0.074	0.069	0.084	0.073	0.076	0.075	0.075	0.077
17-043-6001	Lisle	0.073	0.070	0.071	0.069	0.074	0.071	0.070	0.071
17-049-1001	Effingham	0.062	0.063	0.066	0.070	0.066	0.063	0.066	0.067
17-065-0002	Knight Prairie	0.067	0.064	0.069	0.064	0.068	0.066	0.065	0.067
17-083-1001	Jerseyville	0.062	0.069	-	0.067	0.074	0.067	0.068	0.070
17-085-9991	Stockton	0.063	0.059	0.067	0.063	0.067	0.063	0.063	0.065
17-089-0005	Elgin Larsen Jr. High School	0.073	0.071	0.072	0.069	0.074	0.072	0.070	0.071
17-097-1007	Zion	0.076	0.066	0.074	0.074	0.077	0.072	0.071	0.075
17-111-0001	Cary	0.076	0.070	0.074	0.070	0.073	0.073	0.071	0.072
17-113-2003	Normal	0.070	0.063	0.068	0.064	0.065	0.067	0.065	0.065
17-115-0013	Decatur Illinois EPA Trailer	0.065	0.063	0.069	0.068	0.066	0.065	0.066	0.067
17-117-0002	Nilwood	0.063	0.063	0.066	0.066	0.067	0.064	0.065	0.066

Table B4
Ozone Design Values

AQS ID	City	Fourth High 8-Hour Concentrations (ppm)					Design Values* (ppm)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-119-0120	Alton	0.069	0.067	0.072	0.066	0.073	0.069	0.068	0.070
17-119-1009	Maryville	0.067	0.064	0.075	0.074	0.067	0.068	0.071	0.072
17-119-3007	Wood River	0.069	0.070	0.072	0.067	0.075	0.070	0.069	0.071
17-119-9991	Highland	0.066	0.062	0.071	0.067	0.068	0.066	0.066	0.068
17-143-0024	Peoria Fire Station #8	0.064	0.062	0.069	0.065	0.068	0.065	0.065	0.067
17-143-1001	Peoria Heights	0.070	0.064	0.070	0.066	0.066	0.068	0.066	0.067
17-157-0001	Houston	0.061	0.060	0.065	0.069	0.066	0.062	0.064	0.066
17-161-3002	Rock Island	0.063	0.066	0.067	0.066	0.064	0.065	0.066	0.065
17-163-0010	East St. Louis	0.065	0.064	0.073	0.067	0.073	0.067	0.068	0.071
17-167-0014	Springfield State Fairgrounds	0.067	0.062	0.069	0.069	0.068	0.066	0.066	0.068
17-197-1011	Braidwood	0.067	0.060	0.071	0.068	0.064	0.066	0.066	0.067
17-201-2001	Loves Park	0.067	0.066	0.070	0.064	0.070	0.067	0.066	0.068
Statewide Average		0.069	0.065	0.071	0.068	0.070	0.068	0.068	0.069

*The design value is the three-year average of the fourth high concentration. Design value greater than 0.070 ppm is a violation of the National Ambient Air Quality Standard.

PM_{2.5} FRM and FEM Monitoring Sites



Site ID	Site Name
1. 170190006	Champaign
2. 170191001	Bondville
3. 170310022	Chicago – Washington High School
4. 170310052	Chicago – Mayfair Pump Station
5. 170310057	Chicago – Springfield Pump Station
6. 170310076	Chicago – Com Ed Maint. Bldg.
7. 170310001	Alsip
8. 170310119	Lansing – Kingery near-road
9. 170311016	Lyons Township
10. 170313103	Schiller Park
10. 170313301	Summit
12. 170314007	Des Plaines
13. 170314201	Northbrook
14. 170316005	Cicero
15. 170434002	Naperville
16. 170650002	Knight Prairie
17. 170831001	Jerseyville
18. 170890003	Elgin
19. 170890007	Aurora
20. 171110001	Cary
21. 171132003	Normal
22. 171150013	Decatur
23. 171190024	Granite City – Gateway Medical
24. 171191007	Granite City – 23 rd and Madison
25. 171192009	Alton
26. 171193007	Wood River
27. 171430037	Peoria
28. 171570001	Houston
29. 171613002	Rock Island
30. 171630010	East St. Louis
31. 171670012	Springfield
32. 171971002	Joliet
33. 171971011	Braidwood
34. 172010118	Rockford

Total Over 35 ug/m3	9	
Total Days Over 35 ug/m3	5	

Table B6
PM_{2.5} Highs

AQS ID	City	Total Samples	Samples Greater Than 35 ug/m3			Highest Samples							
			2020	2019	2018	1st	2nd	3rd	4th	5th	6th	7th	8th
17-019-0006	Champaign	112	0	0	0	19.2	17.9	17.3	17.0	16.3	16.2	15.9	15.4
17-019-1001	Bondville	365	0	0	0	22.3	18.9	18.4	18.1	17.7	17.4	17.0	16.1
17-031-0001	Alsip	51	0	0	0	21.5	14.9	14.7	14.5	14.5	14.5	13.1	12.5
17-031-0022	Chicago Washington High School	90	0	0	1	25.5	22.3	19.6	17.3	16.8	16.3	16.1	14.9
17-031-0052	Chicago Mayfair Pump Station	95	1	0	0	36.5	24.0	18.3	17.0	16.3	16.3	15.3	15.2
17-031-0057	Chicago Springfield Pump Station	49	0	0	0	22.4	19.5	16.0	15.8	13.6	13.0	12.6	12.1
17-031-0076	Chicago Com Ed Maintenance	50	0	0	0	14.5	13.5	13.1	13.0	12.7	12.2	12.0	11.7
17-031-0119	Lansing Kingery near-road #1	354	0	0	0	31.8	29.9	28.9	28.2	24.4	23.9	23.8	23.1
17-031-1016	Lyons Township	122	0	0	0	27.0	21.9	19.4	19.3	19.0	19.0	18.0	17.3
17-031-3103	Schiller Park	121	0	0	0	21.4	20.5	20.0	19.8	19.4	19.3	18.6	18.1
17-031-3301	Summit	114	0	0	0	26.6	22.7	21.4	18.9	18.8	17.7	17.2	17.1
17-031-4007	Des Plaines	342	0	0	0	25.8	23.2	21.8	21.0	20.6	18.2	18.0	17.5
17-031-4201	Northbrook	274	0	0	0	17.7	16.7	15.5	15.4	15.1	15.0	15.0	14.9
17-031-6005	Cicero Liberty School	42	0	0	0	21.9	20.7	15.6	14.1	13.6	12.8	12.4	12.3
17-043-4002	Naperville	341	0	0	0	31.3	25.8	23.8	22.8	21.3	21.1	20.9	20.0
17-065-0002	Knight Prairie	348	1	0	0	42.3	27.8	23.9	22.6	19.2	18.0	16.9	16.3
17-083-0117	Jerseyville	354	0	0	0	27.0	21.1	20.7	19.9	19.7	18.0	17.8	16.9
17-089-0003	Elgin McKinley School	94	0	0	0	30.0	25.7	24.0	21.1	19.5	17.8	17.3	14.1
17-089-0007	Aurora	107	0	0	0	32.6	23.9	20.5	19.8	16.3	15.9	15.4	14.8
17-111-0001	Cary	54	0	0	0	24.2	17.1	14.4	14.3	13.0	13.0	12.5	11.9
17-113-2003	Normal	363	0	0	0	26.7	23.7	22.3	19.8	19.4	19.0	18.9	18.9
17-115-0013	Decatur Illinois EPA Trailer	362	0	0	0	19.9	19.8	19.2	19.2	18.5	18.5	17.8	17.6
17-119-0024	Granite City Gateway Medical Center	114	0	0	1	31.1	29.5	23.7	22.4	22.2	21.2	20.1	19.6
17-119-1007	Granite City Fire Station #1	61	0	0	0	23.7	22.3	21.8	17.3	16.5	15.6	15.4	14.8
17-119-2009	Alton SIU Dental Clinic	115	0	0	0	25.5	22.6	22.4	21.6	20.2	20.0	18.3	17.4
17-119-3007	Wood River	116	0	0	0	28.7	26.8	26.1	23.8	22.6	22.5	20.1	18.1
17-143-0037	Peoria	324	1	0	0	40.6	29.6	26.8	24.2	22.7	19.6	19.0	18.2

Table B6
PM_{2.5} Highs

AQS ID	City	Total Samples	Samples Greater Than 35 ug/m3			Highest Samples							
			2020	2019	2018	1st	2nd	3rd	4th	5th	6th	7th	8th
17-157-0001	Houston	357	2	0	0	36.5	35.6	32.4	22.7	20.2	19.0	18.7	18.6
17-161-3002	Rock Island	351	1	0	0	36.1	27.7	22.0	21.4	17.9	17.7	17.3	17.5
17-163-0010	East St. Louis	53	0	0	0	22.2	22.1	21.9	17.6	15.3	15.1	14.6	14.5
17-167-0012	Springfield Agricultural Building	355	0	0	0	32.1	24.5	22.8	21.4	19.8	19.1	18.4	17.6
17-197-1002	Joliet Pershing Elementary	351	2	0	0	62.9	37.2	25.1	23.6	22.7	22.1	22.1	19.6
17-197-1011	Braidwood	349	1	0	0	40.9	26.9	25.3	25.1	20.0	19.6	18.9	18.2
17-201-0118	Rockford Fire Dept.	362	0	1	0	33.2	28.6	23.4	23.1	22.5	22.5	21.3	20.3
Statewide Average						28.9	23.7	21.3	19.8	18.5	17.9	17.3	16.6
Total Over 35 ug/m3			9	1	2								
Total Days Over 35 ug/m3			5	1	2								

Table B7
PM_{2.5} 24-Hour Design Values

AQS ID	City	98th Percentile Concentrations (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-019-0006	Champaign	17.3	19.8	16.8	17.4	15.0	18.0	18.0	16.4
17-019-1001	Bondville	16.1	18.7	17.8	16.7	15.3	17.5	17.7	16.6
17-031-0001	Alsip	14.9	16.0	21.9	20.5	16.9	17.6	19.5	19.8
17-031-0022	Chicago Washington High School	22.3	24.8	27.0	18.3	17.7	24.7	23.4	21.0
17-031-0052	Chicago Mayfair Pump Station	24.0	24.7	25.2	23.3	17.9	24.6	24.4	22.1
17-031-0057	Chicago Springfield Pump Station	22.4	18.6	25.3	20.9	17.5	22.1	21.6	21.2
17-031-0076	Chicago Com Ed Maintenance	14.5	24.9	17.8	23.0	19.0	19.1	21.9	19.9
17-031-0119	Lansing Kingery near-road #1	23.1	21.6	-	-	-	-	-	-
17-031-1016	Lyons Township	19.4	25.8	23.5	23.8	19.9	22.9	24.4	22.4
17-031-3103	Schiller Park	20.0	26.3	25.5	23.8	17.6	23.9	25.2	22.3
17-031-3301	Summit	21.4	19.3	22.5	25.1	17.0	21.1	22.3	21.5
17-031-4007	Des Plaines	18.0	29.0	25.7	22.9	18.9	24.2	25.9	22.5
17-031-4201	Northbrook	15.0	20.7	22.7	20.9	18.4	19.5	21.4	20.7
17-031-6005	Cicero Liberty School	21.9	19.3	22.8	23.6	18.8	21.3	21.9	21.7
17-043-4002	Naperville	20.9	22.8	23.6	22.0	14.8	22.4	22.8	20.1
17-065-0002	Knight Prairie	16.9	17.3	20.6	15.7	16.0	18.3	17.9	17.4
17-083-0117	Jerseyville	16.9	16.9	19.2	19.0	-	17.7	18.4	19.1
17-089-0003	Elgin McKinley School	25.7	24.9	19.5	20.5	15.7	23.4	21.6	18.6
17-089-0007	Aurora	20.5	24.5	21.3	19.8	17.4	22.1	21.9	19.5
17-111-0001	Cary	17.1	18.6	19.0	17.1	14.7	18.2	18.2	16.9
17-113-2003	Normal	18.9	20.6	19.5	18.5	16.3	19.7	19.5	18.1
17-115-0013	Decatur Illinois EPA Trailer	17.6	20.4	22.4	21.6	14.6	20.1	21.5	19.5
17-119-0024	Granite City Gateway Medical Center	23.7	25.0	20.9	16.9	24.7	23.2	20.9	20.8
17-119-1007	Granite City Fire Station #1	22.3	23.8	22.8	21.2	16.2	23.0	22.6	20.1
17-119-2009	Alton SIU Dental Clinic	22.4	19.2	21.8	18.9	20.3	21.1	20.0	20.3

Table B7
PM_{2.5} 24-Hour Design Values

AQS ID	City	98th Percentile Concentrations (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-119-3007	Wood River	26.1	22.7	22.2	17.6	20.7	23.7	20.8	20.2
17-143-0037	Peoria City Office Building	19.6	19.3	20.4	22.4	14.3	19.8	20.7	19.0
17-157-0001	Houston	18.6	16.9	19.1	17.7	18.4	18.2	17.9	19.9
17-161-3002	Rock Island	17.5	20.1	19.4	20.4	17.7	19.0	20.0	19.2
17-163-0010	East St. Louis	22.1	22.9	22.6	18.3	18.4	22.5	21.3	19.8
17-167-0012	Springfield Agricultural Building	17.6	17.9	19.8	20.6	19.1	18.4	19.4	19.8
17-197-1002	Joliet Pershing Elementary	21.0	21.4	20.9	19.6	16.6	21.1	20.6	19.0
17-197-1011	Braidwood	19.1	20.6	19.5	18.5	18.0	19.7	19.5	18.7
17-201-0118	Rockford Fire Department	21.3	23.4	10.6	-	-	18.4	-	-
17-201-0013	Rockford Health Department	-	-	23.0	17.1	14.8	-	-	18.3
Statewide Average		19.9	21.4	21.3	20.1	17.5	20.8	21.0	19.8

*The design value is the three-year average of the 98th percentile concentration. Design value greater than or equal to 35.5 ug/m³ is a violation of the National Ambient Air Quality Standard.

Table B8
PM_{2.5} Annual Design Values

AQS ID	City	Annual Arithmetic Mean Concentrations (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-019-0006	Champaign	7.3	7.5	7.6	7.4	7.6	7.5	7.5	7.5
17-019-1001	Bondville	7.3	7.8	8.0	7.7	7.3	7.7	7.8	7.6
17-031-0001	Alsip	8.6	7.9	9.0	8.7	8.6	8.5	8.5	8.8
17-031-0022	Chicago Washington High School	8.8	10.3	9.6	8.4	8.4	9.6	9.4	8.8
17-031-0052	Chicago Mayfair Pump Station	10.3	9.2	9.8	8.7	8.7	9.8	9.2	9.1
17-031-0057	Chicago Springfield Pump Station	8.3	8.8	9.6	8.9	9.2	8.9	9.1	9.2
17-031-0076	Chicago Com Ed Maintenance	8.3	8.3	9.0	8.4	9.0	8.6	8.6	8.8
17-031-0119	Lansing Kingery near-road #1	10.8	10.8	-	-	-	-	-	-
17-031-3103	Schiller Park	9.9	10.8	11.2	10.3	9.4	10.6	10.8	10.3
17-031-3301	Summit	8.7	9.3	10.2	8.9	9.1	9.4	9.5	9.4
17-031-4007	Des Plaines	8.4	10.3	11.4	9.3	8.9	10.0	10.3	9.9
17-031-4201	Northbrook	7.3	8.5	8.8	8.1	8.0	8.2	8.5	8.3
17-031-6005	Cicero Liberty School	9.3	9.0	10.0	8.6	8.9	9.5	9.2	9.2
17-043-4002	Naperville	9.1	10.3	10.5	8.2	7.8	10.0	9.7	8.8
17-065-0002	Knight Prairie	8.8	8.3	8.9	8.7	7.8	8.6	8.6	8.4
17-083-0117	Jerseyville	7.5	8.0	8.3	8.8	-	7.9	8.4	8.6
17-089-0003	Elgin McKinley School	8.8	8.5	8.7	8.0	7.9	8.7	8.4	8.2
17-089-0007	Aurora	8.4	8.7	9.0	8.1	8.0	8.7	8.6	8.4
17-111-0001	Cary	8.2	7.8	8.2	7.2	7.3	8.1	7.7	7.6
17-113-2003	Normal	8.5	9.2	9.7	8.8	7.6	9.2	9.5	8.7
17-115-0013	Decatur IEPA Trailer	8.6	9.5	10.4	8.7	7.8	9.5	9.5	9.0
17-119-1007	Granite City Fire Station #1	10.1	10.5	11.0	9.6	9.1	10.5	10.4	9.9
17-119-2009	Alton SIU Dental Clinic	9.0	9.1	9.3	8.7	8.8	9.1	9.0	8.9
17-119-3007	Wood River	9.2	9.1	9.2	8.3	8.7	9.2	8.9	8.7
17-143-0037	Peoria City Office Building	8.3	8.0	9.4	8.3	7.6	8.6	8.6	8.5

Table B8
PM_{2.5} Annual Design Values

AQS ID	City	Annual Arithmetic Mean Concentrations (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-157-0001	Houston	8.3	7.7	7.8	9.6	8.0	7.9	8.4	8.4
17-161-3002	Rock Island	8.1	8.6	8.9	7.9	7.2	8.5	8.5	8.0
17-163-0010	East St. Louis	9.5	9.1	10.3	8.8	10.0	9.6	9.4	9.7
17-167-0012	Springfield Agricultural Building	7.6	8.2	9.5	8.6	7.7	8.4	8.8	8.6
17-197-1002	Joliet Pershing Elementary	9.8	9.7	9.8	8.7	8.0	9.8	9.4	8.8
17-197-1011	Braidwood	8.2	8.8	7.9	7.8	7.5	8.3	8.2	7.7
17-201-0118	Rockford Fire Department	9.1	10.3	-	-	-	-	-	-
17-201-0013	Rockford Health Department	-	-	7.7	8.1	7.8	-	-	7.9
Statewide Average		8.7	9.0	9.3	8.5	8.3	9.0	8.9	8.7

*The design value is the three-year average of the annual arithmetic mean concentrations. Design value greater than 12.0 ug/m³ is a violation of the National Ambient Air Quality Standard.

Shaded cells indicate completeness criteria were not met.

PM₁₀ Monitoring Sites



	Site ID	Site Name
1.	170310022	Chicago – Washington High School
2.	170311016	Lyons Township
3.	170314201	Northbrook
4.	171190010	Granite City – 23 rd and Madison

Total Over 150 ug/m3	1	
Total Days Over 150 ug/m3	1	

Table B10
PM₁₀ 24-Hour Highs and Design Values

AQS ID	City	Total Samples	Highest 24-Hour Samples								Samples Greater Than 150 ug/m3			Three-year Exceedance Average*
			1 st	2 nd	3 rd	4 th	5 th	6 th	7 th	8 th	2020	2019	2016	
17-031-0022	Chicago Washington High School	219	96	93	83	83	83	83	76	75	0	0	0	0.0
17-031-1016	Lyons Township	354	159	149	115	102	99	98	95	94	1	0	0	0.3
17-031-4201	Northbrook	58	55	48	45	42	41	38	30	30	0	0	0	0.0
17-119-1007	Granite City Fire Station #1	56	80	73	65	61	54	53	52	50	0	0	0	0.0
Statewide Average			98	91	77	72	69	68	63	62				
Total Over 150 ug/m3											1	0	0	
Total Days Over 150 ug/m3											1	0	0	


*The 24-hour PM₁₀ standard is an exceedance-based standard set at 150 ug/m³. The level is not to be exceeded more than once per year on average over three years. Three-year averages more than one are a violation of the National Ambient Air Quality Standard.

Table B11
PM₁₀ Annual Design Values

AQS ID	City	Annual Arithmetic Mean Concentration (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-031-0022	Chicago Washington High School	32	27	23	24	16	27	25	21
17-031-1016	Lyons Township	37	30	24	25	27	30	26	25
17-031-4201	Northbrook	20	14	14	16	17	16	15	16
17-119-1007	Granite City Fire Station #1	32	35	33	26	28	33	31	29
Statewide Average		30	27	24	23	22	27	24	23

*The annual PM₁₀ standard was revoked in 2007. Previously the standard was a three-year average of the annual means. Concentrations above 50 ug/m³ were a violation of the former National Ambient Air Quality Standard. Currently only the 24-hour PM₁₀ standard is in place (see Table B10).

Carbon Monoxide Monitoring Sites



	Site ID	Site Name
1.	170191001	Bondville
2.	170310119	Lansing - Kingery near-road
3.	170314201	Northbrook
4.	171630010	East St. Louis

Total 1-hour Over 35 ppm	0	Total 8-hour Over 9 ppm	0
Total Days 1-hour Over 35 ppm	0	Total Days 8-hour Over 9 ppm	0

Table B13
Carbon Monoxide Highs


AQS ID	City	Total Hourly Samples	Fourth Highest Daily Samples 1-Hour (ppm)				Fourth Highest Samples 8-Hour (ppm)			
17-019-1001	Bondville	4806	0.21	0.20	0.20	0.16	0.2	0.1	0.1	0.1
17-031-0119	Lansing Kingery near-road #1	5645	1.8	1.7	1.4	1.4	1.4	1.1	1.1	1.0
17-031-4201	Northbrook	6264	0.98	0.92	0.86	0.79	0.7	0.6	0.6	0.5
Statewide Average			1.00	0.94	0.82	0.78	0.77	0.60	0.60	0.53

Table B14
Carbon Monoxide 1-Hour and 8-Hour Design Values

AQS ID	City	1-Hour Samples Greater than 35 (ppm)					8-Hour Samples Greater than 9 (ppm)				
		2020	2019	2018	2017	2016	2020	2019	2018	2017	2016
17-019-1001	Bondville	0	0	0	0	0	0	0	0	0	0
17-031-0119	Lansing Kingery near-road #1	0	0	-	-	-	0	0	-	-	-
17-031-4201	Northbrook	0	0	0	0	0	0	0	0	0	0

*The 1-hour and 8-hour carbon monoxide standard is an exceedance-based standard. The 1-hour standard is set at 35 ppm and is not to be exceeded more than once per year. The 8-hour standard is set at 9 ppm and is not to be exceeded more than once per year. More than one exceedance in a year is a violation of the National Ambient Air Quality Standard.

Sulfur Dioxide Monitoring Sites



Site ID	Site Name
1. 170191001	Bondville
2. 170310076	Chicago – Com Ed Maint. Bldg.
3. 170311601	Lemont
4. 170314201	Northbrook
5. 170990007	Oglesby
6. 171150013	Decatur
7. 171150118	Decatur - Archer Daniel Midlands
8. 171150218	Decatur - Tate & Lyle North
9. 171150318	Decatur - Tate & Lyle South
10. 171170002	Nilwood
11. 171193007	Wood River
12. 171630010	East St. Louis
13. 171790004	Pekin
14. 171850001	Mount Carmel

(74)

Table B16
Sulfur Dioxide Highs


AQS ID	City	Total Hourly Samples	Samples Greater Than 75 ppb			Highest Daily 1-Hour Samples (ppb)				Highest 3- Hour Block Averages (ppb)	
			2020	2019	2018	1st	2nd	3rd	4th	1st	2nd
17-019-1001	Bondville	7492	0	0	0	3.4	3.0	2.7	2.6	2.5	2.3
17-031-0076	Chicago Com Ed Maintenance	8676	0	0	0	17.2	17.2	14.6	14.4	11.5	10.6
17-031-1601	Lemont	8192	0	0	0	8.0	5.5	5.0	4.8	5.9	4.4
17-031-4201	Northbrook	7788	0	0	0	12.6	8.3	6.4	5.4	9.1	4.4
17-099-0007	Oglesby	8278	0	0	0	18.9	8.3	7.8	7.6	6.7	6.6
17-115-0013	Decatur Illinois EPA Trailer	7750	0	0	0	49.5	37.3	32.7	21.6	41.8	24.7
17-115-0117	Decatur ADM	2095	0	0	0	16.3	14.2	11.6	11.3	8.7	7.7
17-115-0217	Decatur Tate & Lyle North	8544	0	0	5	60.6	46.4	39.6	38.8	40.7	37.3
17-115-0317	Decatur Tate & Lyle South	8758	0	0	6	49.0	43.9	41.1	38.5	40.8	31.2
17-117-0002	Nilwood	8401	0	0	0	4.8	3.8	3.5	2.9	4.4	2.7
17-119-3007	Wood River	8364	0	0	0	15.0	9.8	8.2	7.2	8.0	6.1
17-163-0010	East St. Louis	8303	0	0	0	9.9	8.4	8.4	8.3	7.9	6.7
17-179-0004	Pekin	8760	0	0	0	18.5	18.3	15.2	14.3	10.7	10.3
17-185-0001	Mount Carmel	8360	0	0	0	54.0	53.7	50.6	48.9	40.3	35.6
Statewide Average						24.1	19.9	17.7	16.2	17.1	13.6
Total Over 75 ppb			0	0	0						
Total Days Over 75 ppb			0	0	0						

Table B17
Sulfur Dioxide 1-Hour Design Values

AQS ID	City	99th Percentile Concentrations (ppb)					Design Values* (ppb)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-019-1001	Bondville	2.6	3.8	3.3	3.6	3.7	3	4	4
17-031-0076	Chicago Com Ed Maintenance	14.4	10.5	11.0	11.5	9.3	12	11	11
17-031-1601	Lemont	4.8	6.6	6.3	5.3	12.3	6	6	8
17-031-4201	Northbrook	6.1	4.1	3.4	2.5	4.3	5	3	3
17-099-0007	Oglesby	7.6	22.4	27.4	12.5	14.7	19	21	18
17-115-0013	Decatur Illinois EPA Trailer	21.6	23.4	37.0	39.6	54.3	27	33	44
17-115-0117	Decatur ADM	16.3	17.0	20.8	27.8	-	18	22	24
17-115-0217	Decatur Tate & Lyle North	38.8	41.8	83.9	76.6	-	55	67	80
17-115-0317	Decatur Tate & Lyle South	38.5	34.2	89.0	74.3	-	54	66	82
17-117-0002	Nilwood	2.9	4.6	4.5	3.8	5.2	4	4	5
17-119-1010	South Roxana	-	-	-	-	12.9	-	-	-
17-119-3007	Wood River	7.2	9.3	9.7	10.9	24.2	9	10	15
17-143-0024	Peoria Fire Station #8	-	-	-	18.5	27.1	9	-	-
17-163-0010	East St. Louis	8.3	10.6	15.9	8.8	19.1	12	12	15
17-179-0004	Pekin	14.3	17.3	11.8	-	125.8	14	15	69
17-185-0001	Mount Carmel	48.9	30.5	36.8	32.4	42.1	39	33	37
Statewide Average		16.6	16.9	25.8	24.4	27.3	19	22	30

*The design value is the three-year average of the 99th percentile concentration. Design value greater than 75 ppb is a violation of the National Ambient Air Quality Standard.

Nitrogen Dioxide Monitoring Sites



Site ID	Site Name
1. 170310076	Chicago - Com Ed Maintenance
2. 170310216	Chicago - Kennedy near-road
3. 170310116	Lansing - Kingery near-road
4. 170313103	Schiller Park
5. 170314002	Cicero
6. 171170002	Nilwood
7. 171630010	East St. Louis

Total Over 100 ppb	0	
Total Days Over 100 ppb	0	

Table B19
Nitrogen Dioxide Highs

AQS ID	City	Total Hourly Samples	Samples Greater Than 100 ppb			Highest Samples							
			2020	2019	2018	1st	2nd	3rd	4th	5th	6th	7th	8th
17-031-0076	Chicago Com Ed Maintenance	7869	0	0	0	61.1	56.0	50.4	50.1	47.0	45.8	44.6	44.4
17-031-0119	Lansing Kingery near- road #1	8710	0	0	-	58.4	56.2	55.3	53.3	53.0	52.4	51.8	50.4
17-031-0219	Chicago Kennedy near-road #2	8770	0	0	-	61.6	57.2	55.7	53.9	53.4	52.7	51.5	51.1
17-031-3103	Schiller Park	8545	0	0	0	65.6	65.4	62.9	61.9	61.3	59.9	59.5	56.9
17-031-4002	Cicero Cook County Trailer	7053	0	0	0	57.3	56.4	53.2	51.9	51.5	51.0	50.2	50.1
17-117-0002	Nilwood	8676	0	0	0	36.3	34.8	18.4	17.8	16.5	15.9	15.5	15.5
17-163-0010	East St. Louis	8712	0	0	0	46.1	44.3	42.5	42.5	41.5	41.1	40.7	40.6
Statewide Average						55.2	52.9	48.3	47.3	46.3	45.5	44.8	44.1
Total Over 100 ppb			0	0	0								
Total Days Over 100 ppb			0	0	0								

Table B20
Nitrogen Dioxide 1-Hour Design Values

AQS ID	City	98th Percentile Concentrations (ppb)					Design Values* (ppb)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-031-0063	Chicago CTA Building	-	-	-	52.2	58.4	-	-	-
17-031-0076	Chicago Com Ed Maintenance	44.4	46.8	65.9	54.1	60.8	52	56	60
17-031-0119	Lansing Kingery near-road #1	47.8	51.1	-	-	-	49	-	-
17-031-0219	Chicago Kennedy near-road #2	49.9	44.7	-	-	-	47	-	-
17-031-3103	Schiller Park	50.2	54.1	61.0	50.0	56.0	55	55	56
17-031-4002	Cicero Cook County Trailer	49.4	55.7	59.7	55.1	54.7	55	57	57
17-031-4201	Northbrook	-	-	-	-	39.7	-	-	-
17-117-0002	Nilwood	15.5	15.0	15.2	-	-	15	15	-
17-163-0010	East St. Louis	39.1	39.1	38.2	35.9	35.3	39	38	36
Statewide Average		42.3	43.8	48.0	49.5	50.8	45	44	52

*The design value is the three-year average of the 98th percentile concentration. Design value greater than 100 ppb is a violation of the National Ambient Air Quality Standard.

Table B21
Nitrogen Dioxide Annual Design Values

AQS ID	City	Annual Arithmetic Mean Concentrations* (ppb)				
		2020	2019	2018	2017	2016
17-031-0063	Chicago CTA Building	-	-	-	15.75	16.85
17-031-0076	Chicago Com Ed Maintenance	11.33	11.89	15.33	12.86	13.49
17-031-0119	Lansing Kingery near-road #1	16.46	16.64	-	-	-
17-031-0219	Chicago Kennedy near-road #2	14.74	16.37	-	-	-
17-031-3103	Schiller Park	15.19	17.43	17.91	15.79	17.08
17-031-4002	Cicero Cook County Trailer	12.75	14.14	15.89	15.63	14.07
17-031-4201	Northbrook	-	-	-	-	12.10
17-117-0002	Nilwood	2.12	2.37	2.40	-	-
17-163-0010	East St. Louis	8.56	8.82	9.49	8.63	9.12
Statewide Average		11.59	12.52	12.20	13.73	13.79

*The design value is the highest annual average concentration during the most recent two years. Design value greater than 53 ppb is a violation of the National Ambient Air Quality Standard.

Lead Monitoring Sites



	Site ID	Site Name
1.	170310022	Chicago – Washington High School
2.	170310110	Chicago – Perez Elementary
3.	171190010	Granite City – 15 th and Madison

Table B22
Lead Highs

AQS ID	City	Total Sample Days	Highest Monthly Means					Maximum Three-Month Mean
			1st	2nd	3rd	4th	5th	
17-031-0022	Chicago Washington High School	44	0.027	0.025	0.025	0.021	0.016	0.02
17-031-0110	Chicago Perez Elementary	30	0.041	0.026	0.023	0.022	0.016	0.03
17-119-0010	Granite City Air Products	53	0.040	0.036	0.034	0.033	0.028	0.12
Statewide Average			0.036	0.029	0.027	0.025	0.020	0.06

Table B23
Lead Design Values

AQS ID	City	Maximum Three-Month Rolling Mean (ug/m3)					Design Values* (ug/m3)		
		2020	2019	2018	2017	2016	2018-2020	2017-2019	2016-2018
17-031-0022	Chicago Washington High School	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.02
17-031-0110	Chicago Perez Elementary	0.03	0.01	0.01	0.01	0.01	0.03	0.01	0.01
17-031-0113	Chicago ArcelorMittal Steel	-	-	-	-	0.01	-	-	-
17-031-4201	Northbrook	-	-	-	-	0.00	-	-	-
17-089-0113	Geneva Johnson Controls	-	-	-	-	0.05	-	-	-
17-115-0110	Decatur Mueller	-	-	-	0.04	0.04	-	-	-
17-119-0010	Granite City Air Products	0.12	0.15	0.06	0.03	0.02	0.15	0.15	0.06
Statewide Average		0.06	0.06	0.03	0.03	0.02	0.07	0.06	0.03

*The design value is the maximum three-month rolling mean over the latest three-year period. Design value greater than 0.15 ug/m3 is a violation of the National Ambient Air Quality Standard.

Table B24
Filter Analysis Data

AQS ID	City	Total Samples	Highs (ug/m3)		Annual Mean	Total Samples	Highs (ug/m3)		Annual Mean	Total Samples	Highs (ug/m3)		Annual Mean
			1 st	2 nd			1 st	2 nd			1 st	2 nd	
		Cadmium				Chromium				Iron			
17-031-0022	Chicago Washington High School	44	0.005	0.003	0.001	44	0.046	0.032	0.013	44	3.92	2.02	0.66
17-031-0110	Chicago Perez Elementary	30	0.003	0.003	0.001	30	0.041	0.041	0.013	30	0.83	0.81	0.35
17-119-0010	Granite City Air Products	53	0.007	0.006	0.001	53	0.245	0.056	0.018	53	4.95	3.33	0.87
		Manganese				Nickel							
17-031-0022	Chicago Washington High School	44	0.191	0.185	0.054	44	0.017	0.012	0.006				
17-031-0110	Chicago Perez Elementary	30	0.037	0.032	0.014	30	0.012	0.012	0.005				
17-119-0010	Granite City Air Products	53	0.343	0.224	0.058	53	0.041	0.022	0.008				

Table B25
Toxic Compounds

AQS ID	City	Compounds	Highest 24-hour Samples (ppbc)				Annual Average
			1 st	2 nd	3 rd	4 th	
17-031-4201	Northbrook	1,3 Butadiene	0.1	0.1	0.1	0.1	0.05
		Dichloromethane	1.1	1.0	0.9	0.8	0.28
		Chloroform	1.1	1.0	0.8	0.8	0.34
		Carbon Tetrachloride	0.1	0.1	0.1	0.1	0.10
		Tetrachloroethylene	0.2	0.1	0.1	0.1	0.01
		Trichloroethylene	0.0	0.0	0.0	0.0	0.02
		1,2 Dichloropropane	0.0	0.0	0.0	0.0	0.00
		Vinyl Chloride	0.0	0.0	0.0	0.0	0.00
		Benzene	1.4	1.3	1.2	1.1	0.75
		Toluene	4.5	3.6	3.2	3.0	1.53
		Formaldehyde	7.5	4.7	4.2	4.0	2.32
		Acetaldehyde	3.7	3.6	3.4	2.7	1.70
		Acrolein	2.4	2.3	2.3	2.0	0.90
17-031-3103	Schiller Park	1,3 Butadiene	0.3	0.3	0.3	0.3	0.13
		Dichloromethane	0.6	0.6	0.4	0.3	0.18
		Chloroform	0.1	0.1	0.1	0.1	0.03
		Carbon Tetrachloride	0.1	0.1	0.1	0.1	0.10
		Tetrachloroethylene	2.9	2.5	2.4	2.2	1.55
		Trichloroethylene	0.8	0.4	0.4	0.2	0.06
		1,2 Dichloropropane	0.0	0.0	0.0	0.0	0.00
		Vinyl Chloride	0.0	0.0	0.0	0.0	0.00
		Benzene	1.9	1.9	1.9	1.8	1.16
		Toluene	5.9	4.8	4.5	4.4	2.96
		Formaldehyde	11.6	9.6	8.6	8.2	4.98
		Acetaldehyde	23.0	13.2	11.5	8.1	3.59
		Acrolein	2.9	2.5	2.5	2.3	1.26

¹ – Toxic metals data (As, Be, Cd, Cr, Mn, Ni) summarized in Table B24 - Filter Analysis Data

Appendix C: Point Source Emission Inventory Summary

Table C1					
Carbon Monoxide Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
External Fuel Combustion					
Electric Generation	17,065.5	11,188.4	12,253.2	13,628.8	10,592.7
Industrial	5,345.5	5,005.5	4,674.7	4,559.1	4,638.4
Commercial/Institutional	1,493.7	1,345.6	1,433.4	1,445.3	1,497.3
Space Heating	21.3	16.7	17.7	21.4	21.4
Internal Fuel Combustion					
Electric Generation	2,475.6	3,011.5	1,750.4	1,972.8	2,176.3
Industrial	3,552.2	2,847.7	2,648.3	3,188.1	2,962.7
Commercial/Institutional	226.8	187.8	179.0	213.8	294.1
Engine Testing	168.4	165.7	162.1	208.7	212.3
Industrial Processes					
Chemical Manufacturing	1,591.6	1,603.8	1,832.6	1,827.2	1,552.2
Food/Agriculture	1,576.8	1,449.3	1,263.0	1,189.6	1,201.8
Primary Metal Production	13,226.3	10,165.9	9,912.7	12,408.3	21,676.6
Secondary Metal Production	2,492.9	2,105.9	2,103.6	1,906.6	1,893.8
Mineral Products	3,580.7	4,322.5	3,546.7	3,334.4	3,039.8
Petroleum Industry	3,245.9	2,615.6	2,669.7	2,477.7	2,567.2
Paper and Wood Products	0.5	0.5	0.5	0.5	0.2
Rubber and Plastic Products	24.5	21.5	18.5	21.9	21.8
Fabricated Metal Products	214.2	205.8	218.4	191.7	189.2
Oil and Gas Production	241.6	229.5	241.2	244.4	220.5
Miscellaneous Machinery	1.2	0.6	0.6	0.6	0.6
Electrical Equipment	2.0	1.4	1.4	1.4	1.4
Health Services	175.3	171.4	170.9	168.5	164.4
In-Process Fuel Use	403.2	12.0	10.1	112.9	4,567.0
Miscellaneous Manufacturing	37.5	52.2	55.0	59.6	60.3
Organic Solvent Emissions					
Organic Solvent Use	0.2	0.1			
Surface Coating Operations	232.0	235.9	213.4	233.0	237.0
Petroleum Product Storage	0.2	0.2	0.3	0.0	0.2
Bulk Terminals/Plants	26.0	9.9	10.9	17.5	12.6
Printing/Publishing		0.7	0.7	2.1	4.9
Petroleum Marketing/Transport	21.2	21.1	8.4	95.7	22.9
Organic Chemical Storage (large)			0.2	0.0	0.0
Organic Chemical Transportation				3.6	1.0
Organic Solvent Evaporation	9.0	53.6	20.4	39.8	39.8
Solid Waste Disposal					
Government	1,758.0	1,545.9	1,661.5	1,757.6	1,390.7
Commercial/Institutional	40.9	41.0	11.8	11.8	11.8
Industrial	691.7	629.7	663.8	597.5	655.7
Site Remediation	2.2	2.2	2.2	3.3	1.3
Commercial			28.1	15.5	15.2
Institutional				0.1	0.0
Totals	59,944.8	49,267.3	47,785.6	51,961.0	61,945.1

Appendix C: Point Source Emission Inventory Summary

Table C2					
Nitrogen Oxides Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
External Fuel Combustion					
Electric Generation	33,102.0	27,023.2	28,127.4	29,824.7	20,090.9
Industrial	9,217.5	8,425.8	7,863.4	7,392.7	7,419.7
Commercial/Institutional	1,938.0	1,804.4	1,858.3	1,894.3	1,965.9
Space Heating	86.6	66.0	71.9	74.0	59.6
Internal Fuel Combustion					
Electric Generation	2,409.4	3,531.8	2,046.9	2,522.1	2,856.4
Industrial	14,482.6	9,029.6	7,232.8	8,659.5	8,533.5
Commercial/Institutional	541.3	431.2	431.3	471.4	584.4
Engine Testing	563.8	476.6	344.5	327.2	333.5
Industrial Processes					
Chemical Manufacturing	1,552.0	1,363.9	1,452.3	1,468.9	1,437.2
Food/Agriculture	1,504.3	1,346.0	1,299.1	1,137.9	1,191.5
Primary Metal Production	1,329.7	964.5	1,010.2	1,208.4	1,269.9
Secondary Metal Production	667.0	779.6	720.5	629.5	691.6
Mineral Products	5,410.1	7,619.5	6,405.3	6,699.2	6,065.9
Petroleum Industry	4,191.9	3,749.4	3,640.5	3,771.5	3,571.4
Paper and Wood Products	0.9	0.9	0.9	0.9	0.8
Rubber and Plastic Products	26.4	24.1	20.6	27.5	24.0
Fabricated Metal Products	269.8	245.9	266.1	244.2	237.7
Oil and Gas Production	620.6	688.7	691.2	627.8	555.8
Miscellaneous Machinery	0.6	0.8	0.8	0.8	0.8
Electrical Equipment	2.5	1.9	1.9	1.7	1.7
Health Services	6.6	6.6	6.6	7.0	7.0
Textile Products	0.9				
In-Process Fuel Use	190.3	34.0	70.3	165.0	111.8
Miscellaneous Manufacturing	15.7	15.3	18.6	17.9	18.4
Organic Solvent Emissions					
Organic Solvent Use	0.2	0.2			
Surface Coating Operations	420.7	513.0	475.3	473.6	468.3
Petroleum Product Storage			0.2	0.0	0.2
Bulk Terminals/Plants	0.2	0.2	2.9	7.4	12.8
Printing/Publishing	13.3	4.0	0.8	4.0	5.8
Petroleum Marketing/Transport	8.8	0.8	3.5	38.2	11.6
Organic Chemical Storage (large)		8.7	0.2	0.0	0.0
Organic Chemical Transportation				1.5	0.4
Organic Solvent Evaporation	11.3	23.2	15.9	20.2	20.2
Solid Waste Disposal					
Government	592.1	521.6	590.5	574.8	513.4
Commercial/Institutional	13.3	13.3	1.3	1.3	1.3
Industrial	245.7	198.4	201.4	195.2	212.6
Site Remediation	2.8	2.8	2.8	5.8	1.9
Commercial			11.9	11.0	10.9
Institutional				0.1	0.0
Totals	79,438.9	68,915.9	64,888.5	68,507.0	58,289.1

Appendix C: Point Source Emission Inventory Summary

Table C3					
PM ₁₀ Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
External Fuel Combustion					
Electric Generation	4,335.2	3,137.0	2,901.5	4,004.8	2,760.6
Industrial	1,180.1	972.9	734.0	715.8	806.8
Commercial/Institutional	186.6	172.4	179.4	180.8	184.2
Space Heating	3.4	2.8	3.0	3.2	2.6
Internal Fuel Combustion					
Electric Generation	358.4	527.0	291.8	382.1	425.3
Industrial	238.0	218.9	228.7	269.3	258.0
Commercial/Institutional	35.2	23.8	21.9	26.3	29.5
Engine Testing	24.0	20.9	14.7	15.5	15.0
Industrial Processes					
Chemical Manufacturing	1,031.2	978.8	985.4	1,023.5	992.5
Food/Agriculture	5,846.3	5,718.2	5,600.5	5,497.3	5,756.1
Primary Metal Production	872.1	627.0	634.5	882.7	816.4
Secondary Metal Production	955.0	858.6	885.4	869.2	851.6
Mineral Products	4,733.0	4,455.1	4,332.8	4,093.1	3,597.1
Petroleum Industry	1,189.0	1,283.0	1,153.0	1,234.2	1,195.3
Paper and Wood Products	112.7	121.5	130.5	140.9	129.2
Rubber and Plastic Products	168.2	164.6	140.8	162.1	164.7
Fabricated Metal Products	248.4	239.1	258.9	270.0	249.1
Oil and Gas Production	13.4	14.8	14.0	12.0	11.3
Building Construction	0.1	0.0	0.0	0.0	0.0
Miscellaneous Machinery	14.8	15.4	15.2	13.1	13.3
Electrical Equipment	5.1	5.0	5.0	5.1	4.9
Transportation Equipment	0.6	0.1	0.1	0.2	2.2
Health Services	76.9	75.1	79.2	79.0	76.6
Leather and Leather Products	9.7	9.7	11.9	11.9	6.6
Textile Products	0.1	0.0	0.0	0.0	0.0
Type Setting		0.5	0.5	0.5	1.6
Process Cooling	271.6	267.7	237.4	237.7	230.5
In-Process Fuel Use	81.4	0.4	2.9	26.0	42.1
Miscellaneous Manufacturing	19.2	19.0	19.0	51.7	51.9
Organic Solvent Emissions					
Organic Solvent Use	2.9	2.7	23.0	21.4	21.9
Surface Coating Operations	257.4	310.1	250.8	239.9	235.3
Petroleum Product Storage	1.1	1.1	1.1	0.0	0.0
Bulk Terminals/Plants	1.1	2.5	4.1	1.2	1.2
Printing/Publishing	29.3	28.3	29.9	37.6	37.7
Petroleum Marketing/Transport	1.3	1.3	1.0	4.4	4.4
Organic Chemical Storage (large)	5.8	5.7	5.7	6.1	6.1
Dry Cleaning (petroleum based)	0.7	0.7	0.7	7.4	6.2
Organic Solvent Evaporation	5.7	6.3	3.7	10.4	10.4
Solid Waste Disposal					
Government	355.2	351.8	382.7	426.3	410.2
Commercial/Institutional	7.9	7.4	1.3	0.0	0.0
Industrial	92.0	77.1	201.4	86.7	87.8
Site Remediation	14.2	135.5	2.8	13.7	7.2
Commercial			7.2	3.2	3.2
Institutional				0.1	0.0
Totals	22,820.2	20,778.6	19,725.7	21,066.4	19,506.5

Appendix C: Point Source Emission Inventory Summary

Table C4					
Sulfur Dioxide Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
External Fuel Combustion					
Electric Generation	89,806.2	61,147.3	54,066.6	57,192.8	46,507.2
Industrial	19,064.4	16,023.6	13,409.5	12,220.6	11,697.5
Commercial/Institutional	2,582.8	2,405.7	2,486.2	2,606.4	2,515.0
Space Heating	0.6	0.5	0.5	0.5	0.5
Internal Fuel Combustion					
Electric Generation	223.0	271.9	268.5	248.8	294.0
Industrial	62.8	49.0	42.2	70.6	49.8
Commercial/Institutional	24.0	20.1	15.9	16.8	17.3
Engine Testing	8.1	6.7	4.3	5.2	4.3
Industrial Processes					
Chemical Manufacturing	1,330.6	1,000.0	727.9	912.3	850.4
Food/Agriculture	1,192.5	1,097.2	1,440.8	1,436.7	1,301.3
Primary Metal Production	2,046.8	1,413.2	1,426.9	2,533.5	1,624.3
Secondary Metal Production	93.6	92.8	85.7	92.6	73.3
Mineral Products	4,816.4	7,806.9	9,107.2	6,261.1	6,068.8
Petroleum Industry	2,498.1	1,568.3	1,635.0	1,299.7	1,629.0
Paper and Wood Products	0.0	0.0	0.0	0.0	
Rubber and Plastic Products	0.3	0.3	0.2	3.9	0.3
Fabricated Metal Products	15.6	15.1	14.7	12.8	12.7
Oil and Gas Production	1.3	1.2	0.8	0.6	96.7
Miscellaneous Machinery	0.0	0.0	0.0	0.0	0.0
Electrical Equipment	0.0				
Health Services	7.5	7.5	7.5	7.5	7.5
Process Cooling	0.0	0.0	0.0	0.0	0.0
In-Process Fuel Use	175.4	5.7	5.9	61.7	89.9
Miscellaneous Manufacturing	0.5	0.5	0.4	0.4	2.0
Organic Solvent Emissions					
Organic Solvent Use	0.2	0.0			
Surface Coating Operations	9.6	4.5	4.5	4.9	4.8
Petroleum Product Storage	8.3	0.9	8.3	8.3	8.3
Bulk Terminals/Plants				0.5	0.5
Printing/Publishing	0.8	0.8	0.5	0.0	0.0
Petroleum Marketing/Transport	75.3	0.0	0.0	2.5	1.2
Organic Chemical Storage (large)		0.1	0.1	0.5	0.5
Organic Chemical Transportation	0.1	0.3	1.6		0.0
Organic Solvent Evaporation	3.5	0.7	0.6	0.9	0.9
Solid Waste Disposal					
Government	949.8	729.9	1,063.8	900.8	714.7
Commercial/Institutional	2.6	2.5	1.5	1.5	1.5
Industrial	342.5	371.8	365.7	218.4	232.8
Site Remediation				1.8	1.0
Commercial				0.7	0.7
Institutional				0.0	0.0
Totals	125,421.1	94,095.4	86,245.4	86,125.6	73,808.5

Appendix C: Point Source Emission Inventory Summary

Table C5					
Volatile Organic Material Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
External Fuel Combustion					
Electric Generation	1,095.4	973.2	1,111.1	1,128.9	701.3
Industrial	321.4	338.8	314.9	303.9	306.2
Commercial/Institutional	86.7	78.9	83.7	85.5	89.7
Space Heating	4.6	3.5	3.8	3.9	3.2
Internal Fuel Combustion					
Electric Generation	387.6	528.2	352.7	172.2	219.6
Industrial	793.6	602.8	519.0	684.8	638.7
Commercial/Institutional	35.1	36.6	36.2	45.6	59.0
Engine Testing	39.1	35.3	45.0	56.7	57.4
Industrial Processes					
Chemical Manufacturing	6,261.4	5,752.3	5,769.7	5,679.5	5,658.5
Food/Agriculture	9,461.8	8,917.4	9,316.2	9,432.5	9,718.2
Primary Metal Production	287.8	141.1	146.8	163.6	256.1
Secondary Metal Production	697.4	672.8	725.7	760.1	776.7
Mineral Products	1,163.9	1,257.7	1,100.6	999.7	939.8
Petroleum Industry	1,987.0	1,833.9	1,979.2	1,748.7	1,775.0
Paper and Wood Products	78.4	64.4	59.5	68.3	64.0
Rubber and Plastic Products	1,839.3	1,646.5	1,670.1	1,603.5	1,471.9
Fabricated Metal Products	689.8	790.5	648.2	667.7	710.6
Oil and Gas Production	327.4	351.3	303.7	288.9	269.4
Miscellaneous Machinery	83.4	83.5	74.2	31.1	27.8
Electrical Equipment	38.9	65.7	68.0	65.2	63.8
Transportation Equipment	18.5	18.5	18.5	18.2	18.2
Health Services	12.6	11.8	10.6	5.8	3.9
Photographic Film Manufacturing		1.7	1.7	0.8	
Leather and Leather Products	16.9	16.9	17.9	17.9	17.2
Textile Products	2.3	2.3	2.3	2.3	2.3
Process Cooling	78.9	80.7	80.7	80.7	79.5
In-Process Fuel Use	9.6	6.7	6.7	10.6	11.5
Miscellaneous Manufacturing	139.3	136.2	104.7	67.4	67.5
Organic Solvent Emissions					
Organic Solvent Use	394.	449.4	472.5	502.0	512.9
Surface Coating Operations	6,879.4	6,264.5	6,138.0	6,064.1	5,656.8
Petroleum Product Storage	2,524.1	2,482.5	2,517.0	2,492.5	2,368.0
Bulk Terminals/Plants	1,162.7	1,012.2	1,015.6	1,052.0	895.5
Printing/Publishing	3,081.6	2,451.1	2,467.7	2,382.2	1,947.2
Petroleum Marketing/Transport	434.5	450.4	354.7	358.5	344.3
Organic Chemical Storage (large)	705.5	514.01	578.7	775.3	861.7
Organic Chemical Transportation	102.5	101.4	60.6	41.6	48.9
Dry Cleaning (petroleum based)	374.8	318.0	283.5	280.8	232.7
Aerosol Can Filling					170.1
Organic Chemical Storage (small)	0.2	0.2	0.2		

Appendix C: Point Source Emission Inventory Summary

Table C5					
Volatile Organic Material Point Source Emission Distribution (Tons/Year)					
Category	2016	2017	2018	2019	2020
Organic Solvent Evaporation	416.3	410.9	372.0	354.5	341.4
Solid Waste Disposal					
Government	359.4	413.9	514.5	407.5	403.9
Commercial/Institutional	3.8	3.8	2.9	2.9	2.9
Industrial	58.2	54.6	61.3	60.3	60.3
Site Remediation	142.2	150.3	139.8	97.5	90.9
Commercial				3.9	3.7
Institutional				0.0	0.0
Totals	42,884.5	39,768.0	39,785.1	39,070.1	37,948.2

Appendix C: Point Source Emission Inventory Summary

Table C6					
2020 Estimated County Stationary Point Source Emissions (Tons/Year)					
County	Carbon Monoxide	Nitrogen Oxides	PM ₁₀	Sulfur Dioxide	Volatile Organic Material
Adams	311.8	212.1	249.4	506.7	1,083.5
Alexander	27.2	32.0	40.8	0.4	648.5
Bond	10.2	11.3	11.3	0.7	24.2
Boone	47.4	58.3	57.6	1.0	437.4
Brown	0.0	0.0	2.8	0.0	0.0
Bureau	14.8	28.2	59.8	0.3	34.6
Calhoun	0.6	0.7	5.2	0.0	0.1
Carroll	27.9	27.8	29.6	1.1	11.4
Cass	40.7	46.5	45.2	29.9	13.8
Champaign	296.9	599.4	195.8	102.5	401.2
Christian	316.9	1,152.7	126.0	1,533.8	295.0
Clark	40.7	4.9	61.7	1.4	205.1
Clay	4.0	6.1	15.6	0.1	87.5
Clinton	265.8	807.7	68.2	287.0	67.6
Coles	85.5	81.2	69.2	6.7	150.2
Cook	11,343.8	4,461.5	2,303.0	2,006.3	6,152.6
Crawford	989.0	1,537.1	490.1	4,316.0	817.0
Cumberland	13.6	3.2	17.0	1.0	41.7
DeKalb	137.1	104.9	80.1	38.1	125.5
DeWitt	71.9	60.2	106.7	15.0	188.8
Douglas	804.9	1,448.2	100.2	0.8	455.6
DuPage	690.2	798.7	265.4	38.1	1,103.9
Edgar	15.2	71.1	67.3	0.1	82.5
Edwards	1.3	3.9	10.2	0.0	8.7
Effingham	10.0	24.2	53.7	1.7	264.6
Fayette	61.4	283.0	16.4	247.6	25.7
Ford	82.0	162.4	182.3	5.0	768.0
Franklin	10.0	5.5	28.1	0.0	17.9
Fulton	8.9	8.2	12.5	0.0	13.1
Gallatin	0.0	0.0	7.4	0.0	0.0
Greene	0.1		19.3		0.2
Grundy	630.7	990.2	182.9	40.9	535.2
Hamilton	0.3	0.5	63.9	0.0	0.9
Hancock	15.3	3.2	51.5	0.4	9.2
Hardin	1.6	1.9	12.5	0.0	1.9
Henderson			30.1		
Henry	642.7	1,180.1	161.2	17.8	331.3
Iroquois	56.1	29.2	136.0	4.3	455.8
Jackson	370.7	360.4	63.9	239.4	60.6
Jasper	2,246.8	1,814.6	120.1	5,004.5	120.0
Jefferson	58.1	65.6	30.3	0.5	273.9
Jersey	0.1		6.1		10.3
Jo Daviess	202.3	397.7	131.5	21.5	53.8
Johnson	25.1	24.0	7.8	220.0	5.9
Kane	353.2	359.9	209.8	24.9	880.1
Kankakee	1,036.9	705.2	183.7	54.4	755.1
Kendall	278.1	618.8	263.2	25.8	95.4
Knox	25.9	22.8	55.5	1.9	76.3
Lake	1,613.4	1,538.0	540.0	925.2	467.9
La Salle	1,418.0	2,461.3	968.5	437.1	1,140.8
Lawrence	8.7	5.1	14.6	0.6	36.1
Lee	214.7	312.2	116.6	24.4	263.7

Appendix C: Point Source Emission Inventory Summary

Table C6					
2020 Estimated County Stationary Point Source Emissions (Tons/Year)					
County	Carbon Monoxide	Nitrogen Oxides	PM ₁₀	Sulfur Dioxide	Volatile Organic Material
Livingston	497.0	274.2	141.0	98.5	274.4
Logan	28.4	38.5	69.6	427.8	8.6
McDonough	37.4	72.6	24.3	0.8	68.8
McHenry	197.3	234.0	109.2	5.5	254.7
McLean	261.9	270.9	158.1	15.4	699.4
Macon	1,210.0	5,033.0	2,047.3	10,873.7	4,438.4
Macoupin	6.2	6.7	35.9	0.0	5.0
Madison	19,497.6	3,260.0	1,240.8	2,731.3	2,382.9
Marion	22.8	37.6	35.0	2.2	595.1
Marshall	31.2	74.6	135.1	239.0	422.4
Mason	2.1	2.6	43.2	0.0	34.6
Massac	3,437.1	3,670.6	595.6	10,655.1	120.6
Menard	15.1	3.3	16.4	0.0	26.4
Mercer	0.4	0.5	17.0	0.0	14.3
Monroe	2.8	4.1	12.0	0.1	8.2
Montgomery	32.9	8.6	47.6	4.2	28.5
Morgan	64.8	191.3	48.2	23.3	27.5
Moultrie	3.1	9.3	26.9	0.0	47.0
Ogle	544.0	377.4	326.8	240.8	489.2
Peoria	1,646.7	3,239.0	408.0	6,930.4	822.0
Perry	21.6	49.1	76.1	0.5	41.9
Piatt	95.2	927.2	44.8	0.1	58.9
Pike	88.3	118.4	76.9	2.1	41.5
Pope					
Pulaski	75.9	12.9	47.5	4.1	7.7
Putnam	182.7	213.9	203.1	691.5	171.4
Randolph	970.8	2,838.6	123.4	2,280.2	215.0
Richland	25.1	31.6	15.1	0.2	10.7
Rock Island	304.7	340.4	162.0	24.4	563.5
St. Clair	326.4	287.7	235.2	102.4	484.4
Saline	75.0	51.6	21.6	3.5	8.4
Sangamon	658.1	1,198.0	215.3	1,351.3	164.3
Schuyler	0.0	0.0	9.3	0.0	20.5
Scott	41.8	40.8	51.0	6.6	3.7
Shelby	44.2	127.8	63.2	2.2	57.4
Stark			18.0		6.4
Stephenson	73.1	103.0	105.8	8.4	88.2
Tazewell	490.2	2,091.8	756.7	1,958.3	686.3
Union	37.0	45.9	33.0	687.5	1.8
Vermilion	330.9	444.6	188.5	11.1	1,669.4
Wabash	8.9	2.9	31.5	0.0	5.2
Warren	65.5	36.0	64.9	149.4	14.1
Washington	329.5	4,039.0	1,390.7	10,429.7	130.8
Wayne	44.1	80.0	8.1	4.1	14.2
White	47.5	24.8	2.8	3.0	39.4
Whiteside	630.6	183.7	141.5	22.3	70.3
Will	3,013.1	3,530.5	1,262.7	1,567.3	2,574.3
Williamson	1,090.3	1,245.6	133.0	5,846.4	226.6
Winnebago	400.6	469.9	295.3	217.0	625.5
Woodford	6.7	12.8	43.9	1.7	78.4

Appendix C: Point Source Emission Inventory Summary

Table C7					
Annual Source Estimated Emissions Trends (Tons)					
Year	Carbon Monoxide	Nitrogen Oxides	PM₁₀	Sulfur Dioxide	Volatile Organic Material
1981	240,421	826,427		1,577,992	270,814
1982	163,704	693,054		1,404,040	233,951
1983	144,622	759,453		1,363,292	207,405
1984	110,922	746,367		1,435,066	197,418
1985	107,876	715,556		1,406,300	191,070
1986	109,777	676,181		1,400,761	180,148
1987	98,213	644,511		1,379,407	176,406
1988	127,758	653,521		1,393,628	165,792
1989	132,214	610,214		1,254,474	193,499
1990	134,744	623,466		1,272,445	170,378
1991	148,667	619,161		1,239,690	154,008
1992	129,054	610,214	181,775	1,228,949	156,867
1993	130,097	556,460	113,482	1,170,549	152,288
1994	127,848	555,893	50,730	1,158,555	140,492
1995	127,661	505,966	48,839	1,273,786	141,381
1996	130,040	495,267	43,950	1,183,278	139,445
1997	117,046	510,729	41,078	1,197,404	136,541
1998	108,117	509,676	43,392	1,196,461	134,924
1999	120,906	421,993	40,598	1,085,828	99,121
2000	122,702	424,609	36,885	1,070,058	101,147
2001	96,970	358,263	34,233	653,797	95,221
2002	99,173	301,216	30,422	531,343	90,014
2003	88,367	289,921	41,589	512,321	89,579
2004	80,479	248,245	42,402	507,142	84,080
2005	83,671	238,026	40,359	522,677	75,690
2006	89,717	219,200	37,979	487,588	70,858
2007	80,969	205,602	34,847	429,976	59,021
2008	80,628	203,014	34,474	406,905	57,135
2009	78,720	198,178	32,551	375,807	54,668
2010	65,797	138,344	30,931	304,709	49,975
2011	78,283	143,035	29,796	295,658	48,323
2012	76,255	131,326	28,624	276,412	46,957
2013	64,915	109,308	25,744	211,873	45,430
2014	67,921	109,444	24,942	200,350	44,610
2015	66,072	99,753	23,959	182,200	42,345
2016	59,945	79,439	22,820	125,421	42,885
2017	49,267	68,916	20,779	94,095	39,768
2018	47,786	64,889	19,726	86,245	39,785
2019	51,961	68,507	21,066	86,126	39,070
2020	61,945	58,289	19,507	73,809	37,948

Appendix C: Point Source Emission Inventory Summary

Table C8

Annual Source Reported Emissions Trends (Tons)

Year	Carbon Monoxide	Nitrogen Oxides	PM ₁₀	Sulfur Dioxide	Volatile Organic Material
1992	112,403	381,938	49,377	1,045,113	143,853
1993	113,781	418,209	36,737	1,001,123	108,847
1994	116,192	404,486	34,086	967,213	108,897
1995	160,256	366,978	31,491	814,229	103,144
1996	84,258	407,683	30,850	914,295	87,271
1997	71,408	404,289	25,648	974,232	76,350
1998	79,147	377,191	31,828	964,262	77,952
1999	91,153	360,850	27,663	863,759	71,514
2000	90,315	329,141	30,482	620,592	71,063
2001	83,453	291,778	28,929	531,504	62,647
2002	83,795	261,202	26,900	498,754	70,703
2003	75,511	230,068	29,939	507,338	63,495
2004	77,847	229,127	31,896	521,808	64,594
2005	85,892	215,366	30,535	486,534	62,251
2006	77,099	200,832	29,367	429,573	53,791
2007	77,211	198,073	28,784	406,405	50,933
2008	75,183	193,637	28,194	376,627	49,112
2009	62,285	134,274	25,988	305,297	41,839
2010	75,277	139,508	25,993	297,254	44,245
2011	73,586	129,058	25,209	272,747	42,430
2012	64,253	109,298	22,631	220,143	42,735
2013	65,879	107,877	21,549	201,509	41,276
2014	65,865	99,230	21,962	182,337	40,767
2015	57,688	80,469	19,557	136,749	40,039
2016	46,864	68,441	17,560	99,907	37,593
2017	46,747	64,673	17,209	86,446	37,206
2018	50,727	68,632	18,316	87,437	37,265
2019	61,586	62,595	16,582	78,506	36,723

Appendix D: Website Links

Illinois EPA's Website Information

To access the online version of the Annual Air Quality Report, various pollutant averages and exceedances, the monitoring network plan and emission trends:

- <https://www2.illinois.gov/epa/topics/air-quality/Pages/default.aspx>

Air Quality Index Information

To view current Air Quality Index numbers and forecasts across the country:

- <http://www.airnow.gov>

To sign up for air quality information such as forecasts and pollution alerts:

- <http://www.illinois.enviroflash.info/signup.cfm>

EnviroFlash on Twitter:

- <http://www.illinois.enviroflash.info/EnviroFlashTwitter.cfm>

Monitoring Data Access Information

To access yearly Air Quality Index summaries, air quality statistics and monitoring concentrations:

- <https://www.epa.gov/outdoor-air-quality-data>

To access status and trends of key air pollutants:

- <https://www.epa.gov/air-trends>

To access historical Design Values (statistic to compare to the National Ambient Air Quality Standards):

- <https://www.epa.gov/air-trends/air-quality-design-values>

Nonattainment Areas and Designations (regions in violation of the various National Ambient Air Quality Standards):

- <http://www.epa.gov/green-book>

Other

- Ambient Monitoring Technology Information Center: <https://www.epa.gov/amtic>
- Toxic Release Inventory Search: <http://www.epa.gov/enviro/tri-search>
- Toxic Release Inventory Data and Tools: <https://www.epa.gov/toxics-release-inventory-tri-program/tri-data-and-tools>