



HEALTH ADVISORY FOR 6:2 FLUOROTELOMER SULFONIC ACID (6:2 FTS) CHEMICAL ABSTRACT SERVICES REGISTRY NUMBER (CASRN) 27619-97-2

Prepared by:
Office of Toxicity Assessment
Illinois Environmental Protection Agency
November 6, 2025

REASON FOR ACTION

As a result of Per- and Polyfluoroalkyl Substances (PFAS) sampling in community water supplies (CWS) within the state, 6:2 Fluorotelomer sulfonic acid (6:2 FTS) has been confirmed in a well at a CWS. In accordance with 35 Illinois Administrative Code 620.605(a), the Illinois EPA is issuing a health advisory for 6:2 FTS. Section 620.605(a) directs the Illinois EPA to issue a health advisory for a chemical substance if all of the following conditions are met:

- 1) A community water supply well is sampled, and a substance is detected and confirmed by resampling;
- 2) There is no standard under Section 620.410 for such chemical substance; and
- 3) The chemical substance is toxic or harmful to human health according to the procedures of Appendix A, B, or C.

The health advisory guidance level for 6:2 FTS is 0.00077 milligrams per liter (mg/L), or 770 nanograms per liter (ng/L) or parts per trillion (ppt).



The health advisory will be published in the *Environmental Register* (publication of the Illinois Pollution Control Board), and placed at the website:

<https://pcb.illinois.gov/Resources/News>

The health advisory will also be placed on Illinois EPA's website at:

<https://epa.illinois.gov/topics/water-quality/pfas/pfas-healthadvisory.html>

PURPOSE OF A HEALTH ADVISORY

In accordance with 35 Ill. Adm. Code 620.601, the purpose of a health advisory is to provide guidance levels that, in the absence of an applicable groundwater quality standard under Section 620.410, must be considered by Illinois EPA in: 1) establishing groundwater cleanup or action levels whenever there is a release or substantial threat of a release of a hazardous substance, pesticide, or another contaminant that represents a significant hazard to public health or the environment; 2) determining whether a community water supply is taking its raw water from a site or source consistent with regulatory requirements; and 3) developing Illinois Pollution Control Board (Board) rulemaking proposals for new or revised numerical standards.

Health advisories serve as informal technical guidance, intended to provide information about contaminant exposures and potential public health impacts. The guidance levels represent concentrations in drinking water at which no adverse health effects are expected to occur. Guidance levels are not enforceable or intended to be used as drinking water standards, also known as maximum contaminant levels (MCLs).

HEALTH ADVISORY GUIDANCE LEVEL FOR 6:2 FTS

Through issuance of this Health Advisory, Illinois EPA is providing public notice of its guidance level for 6:2 FTS in drinking water. For non-carcinogenic health effects, the guidance level is 0.00077 milligrams per liter (mg/L), or 770 nanograms per liter (ng/L) or parts per trillion (ppt).

Section 620.605 prescribes the methods for developing health advisories for carcinogens and non-carcinogens. 6:2 FTS does not meet the definition of a "carcinogen" as defined in Section 620.110; therefore, the method for developing a health advisory for non-carcinogens was used. Briefly, this method specifies that the United States Environmental Protection Agency (U.S. EPA) MCL or maximum contaminant level goal (MCLG) is the guidance level, if available, or the human threshold toxicant advisory concentration (HTTAC) must be determined using the procedures contained in Appendix A of Section 620. U.S. EPA has not published an MCL or MCLG for 6:2 FTS; therefore, Illinois EPA used the Appendix A procedures to calculate an HTTAC for 6:2 FTS.

Appendix A specifies, in prescribed order, the toxicological data to be used in developing guidance levels. To determine appropriate toxicological data in accordance with nationally accepted guidelines, pursuant to the Illinois Groundwater Protection Act (415 ILCS 55-8(a)), Illinois EPA relied upon U.S. EPA guidance titled, “Tier 3 Toxicity Value White Paper” (paper), dated May 16, 2013, prepared by the U.S. EPA Office of Solid Waste and Emergency Response (OSWER) Human Health Regional Risk Assessors Forum. The paper lists a hierarchy of sources to be used when determining an appropriate toxicological value for use in human health assessments. The hierarchy for selection of toxicity values is as follows:

Tier 1: U.S. EPA Integrated Risk Information System (IRIS).

Tier 2: U.S. EPA Provisional Peer-Reviewed Toxicity Values (PPRTVs).

Tier 3: In the order in which they are presented:

- 1) The U.S. Health and Human Services Agency for Toxic Substances and Disease Registry (ATSDR) Dose Minimal Risk Levels (dose MRLs).
- 2) California EPA, Office of Environmental Health Hazard Assessment (OEHHA).
- 3) PPRTV “Appendix” Values.
- 4) Health Effects Assessment Summary Table (HEAST).

The paper also references peer-reviewed toxicity values developed by other federal programs to calculate provisional drinking water health advisory levels as a Tier 3 source. In 2022, U.S. EPA placed Office of Water PFAS toxicity values above California EPA’s OEHHA toxicity values within the Tier 3 hierarchy and more recently added peer-reviewed toxicity values derived by U.S. EPA Office of Research and Development (ORD) and Wisconsin.

In December 2024, ORD, under the Health and Environmental Risk Assessment (HERA) National Research Program published a peer-reviewed toxicological assessment titled, “ORD Human Health Toxicity Value for 6:2 Fluorotelomer Sulfonic Acid (CASRN 27619-97-2 |DTXSID6067331).” ORD’s toxicological assessment recommends a chronic oral reference dose (RfD) equal to 0.0002 mg/kg-day. The value is based on a critical effect of decreased immune response in adult male mice from a study by Bohannon et al. titled, “Next-generation PFAS 6:2 Fluorotelomer Sulfonate Reduces Plaque Formation in Exposed White-Footed Mice,” published in 2023. Benchmark dose modeling was used to find the point of departure (POD) for 6:2 FTS of 1.51 mg/kg-day. The human equivalent dose was

calculated using a dosimetric adjustment factor (DAF) to derive the point of departure human equivalent dose (POD_{HED}) of 0.200 mg/kg-day.

A total composite uncertainty factor (UF) of 1,000 (UF of 3 to account for toxicodynamic differences between humans and animals, UF of 10 to account for intraspecies variability, UF of 1 to account for the POD being a BMDL, UF of 3 to account for extrapolation from subchronic to chronic, and UF of 10 to account for database uncertainties) was applied to the POD_{HED}.

The overall RfD for 6:2 FTS was calculated by dividing the POD_{HED} by the composite uncertainty factor.

$$RfD = \frac{POD_{HED}}{UF}$$
$$RfD = \frac{0.200 \text{ mg/kg-day}}{1,000}$$
$$RfD = 0.0002 \text{ mg/kg-day}$$

Using the RfD of 0.0002 (2E-4) mg/kg-day, and the procedures outlined in Section 620. Appendix A, the recommended guidance level for drinking water is 0.00077 milligrams per liter (mg/L), or 770 nanograms per liter (ng/L) or parts per trillion (ppt).


CHEMICAL CHARACTERISTICS AND POTENTIAL ADVERSE HEALTH EFFECTS

General Description of 6:2 FTS


6:2 Fluorotelomer sulfonic acid, also known as 1H,1H, 2H, 2H-Perfluorooctane sulfonic acid or 6:2 FTS, is a synthetic chemical which is part of a larger class of chemicals referred to as per- and polyfluoroalkyl substances. PFAS have been manufactured since the middle 20th Century and are known for their chemical and physical properties that impart oil and water repellency, temperature resistance, and friction reduction to a wide range of products, including, but not limited to, textile coatings, paper products, food wrappers, cosmetic and personal care products, non-stick cookware and fire-fighting foams. PFAS are also used in the semiconductor, aerospace, oil production and mining, and metal plating industries, to name a few. PFAS enter the environment through industrial manufacturing and the use and disposal of PFAS-containing products. The chemical and physical properties of 6:2 FTS make it mobile, persistent, and bioaccumulative, meaning fish and other animals may accumulate 6:2 FTS in animal tissue when their food sources are contaminated with 6:2 FTS. 6:2 FTS is known to be persistent in the environment.

Manufacturers designed 6:2 FTS as a replacement product for perfluorooctanesulfonic acid (PFOS) in fire-fighting foam and other uses.

Structural Identifier



Chemical Identifier



Potential Adverse Health Effects of 6:2 FTS

Epidemiological studies on human health effects from exposure to 6:2 FTS are limited in their ability to draw conclusions on the associations between health effects and exposure.

Information regarding health effects of 6:2 FTS are primarily derived from animal studies, via the ingestion, or oral exposure, route. Laboratory studies observed the following effects in animals exposed to 6:2 FTS:

- Increased liver weight
- Reduced body weight
- Decreased immune response

Carcinogenic Potential

Section 620.110. defines a carcinogen as a contaminant that is classified as: 1) a Category A1 or A2 Carcinogen by the American Conference of Governmental Industrial Hygienists (ACGIH); 2) a Category 1 or 2A/2B Carcinogen by the World Health Organization's International Agency for Research on Cancer (IARC); 3) a "Human Carcinogen" or "Anticipated Human Carcinogen" by the United States Department of Health and Human

Service National Toxicological Program (NTP); or 4) a Category A or B1/B2 Carcinogen by the U.S. EPA in IRIS or a Final Rule issued in a Federal Register notice by the USEPA. 6:2 FTS is not classified as a carcinogen by any of the above sources.

**ATTACHMENT TO HEALTH ADVISORY
FOR
6:2 FLUOROTELOMER SULFONIC ACID (6:2 FTS)
CASRN 27619-97-2**

OVERVIEW OF KEY STUDIES

For information regarding the studies used by U.S. EPA's ORD for the derivation of its 6:2 FTS RfD, refer to "ORD Human Health Toxicity Value for 6:2 Fluorotelomer Sulfonic Acid," located at: <https://assessments.epa.gov/risk/document/&deid%3D363720>.

DERIVATION OF THE HEALTH ADVISORY GUIDANCE LEVEL FOR 6:2 FTS

The first step in the derivation of a health advisory guidance level is to determine whether the chemical substance presents a carcinogenic risk to humans. 6:2 FTS does not meet the definition of a carcinogen as specified in Part 620. Therefore, the guidance level will be based on noncarcinogenic effects of this chemical.

In deriving a guidance level to protect against a health effect for which there is a threshold dose below which no damage occurs (i.e., noncarcinogen effects), Section 620.605 specifies that U.S. EPA's MCLG, if available, is the guidance level. U.S. EPA has not published a MCLG for 6:2 FTS; therefore, Illinois EPA must calculate the HTTAC as the guidance level, using the procedures specified in Appendix A of Section 620.

Appendix A specifies in subsection (a) that the HTTAC is calculated as follows:

$$HTTAC = \frac{RSC \cdot ADE}{W}$$

Where:

- HTTAC = Human threshold toxicant advisory concentration in milligrams per liter (mg/L).
- RSC = Relative source contribution, the relative contribution of the amount of exposure to a chemical via ingestion of drinking water when compared to total exposure to that chemical from all sources. Valid chemical-specific data shall be used if available. If valid chemical-specific data are not available, a value of 20% (= 0.20) must be used.
- ADE = Acceptable daily exposure of a chemical in milligrams per day (mg/d) as determined in accordance with Appendix A, subsection (b).

W = Per capita daily water consumption for a child (0-6 years of age), equal to 0.78 liters per day (L/d).

Subsection (b) of Appendix A specifies that the ADE be calculated using, in specified order: a U.S. EPA verified RfD (an estimate of a daily exposure to a chemical which is expected to be without adverse health effects for humans for a lifetime of exposure in units of mg/kg-day); a NOAEL which has been identified as a result of human exposures; a LOAEL which has been identified as a result of human exposures; a NOAEL which has been determined from studies with laboratory animals; and a LOAEL which has been determined from studies with laboratory animals.

Illinois EPA selected the U.S. EPA ORD recommended RfD of 0.0002 (2E-4) mg/kg-day, as the verified RfD for use in calculating the ADE. The ADE equals the product of multiplying the toxicity value by 15 kilograms (kg), which is the assumed average body weight of a child 0-6 years of age per Section 620:

$$ADE = 0.0002 \text{ mg/kg-day} \cdot 15 \text{ kg} = 0.003 \text{ mg/day}$$

The next step in the development of the HTTAC is the evaluation of chemical-specific RSC data available for the chemical. Illinois EPA evaluated data from ATSDR, U.S. EPA Office of Water, and values developed by other states. There is little scientific consensus regarding the contribution of drinking water to the total amount of PFAS exposure to humans. Humans are exposed to 6:2 FTS through a variety of media, including, but not limited to air emissions, ingestion of fish or other animals exposed to 6:2 FTS, dermal exposure and incidental exposure from 6:2 FTS-containing consumer products, much of which varies on a site-specific basis. Due to this lack of consensus, Illinois EPA elected to use the conservative default value of 20% (0.20) for its HTTAC calculation.

The HTTAC is calculated by the product of the RSC and the ADE, divided by the per capita daily water ingestion rate, specified in Appendix A as equal to 0.78 L/day:

$$HTTAC \text{ (mg/L)} = \frac{0.20 \cdot 0.003 \text{ mg/day}}{0.78 \text{ L/day}}$$

$$HTTAC \text{ (mg/L)} = \frac{0.0006 \text{ mg/day}}{0.78 \text{ L/day}}$$

$$HTTAC = 0.00077 \text{ mg/L}$$

or:

770 ng/L or ppt

The final step in ensuring a calculated guidance level is appropriate is to compare the guidance level to the chemical's practical quantitation limit (PQL), or minimum reporting level (MRL). U.S. EPA's Method 533 for analyses of PFAS drinking water samples shows the 6:2 FTS MRL is 5 ng/L, which is below the calculated guidance level of 770 ng/L. Therefore, the guidance level is appropriate.

REFERENCES

Bohannon, ME; Narizzano, AM; Guigni, BA; East, AG; Quinn, MJ. (2023). Next-generation PFAS 6:2 fluorotelomer sulfonate reduces plaque formation in exposed white-footed mice. *Toxicol Sci* 192: 97-105. <http://dx.doi.org/10.1093/toxsci/kfad006>.

IGA (Illinois General Assembly). Illinois Groundwater Protection Act (IGPA). 415 ILCS 55. Available at: <https://www.ilga.gov/legislation/ilcs/ilcs3.asp?ActID=1595&ChapterID=36>

PCB (Pollution Control Board). Title 35: Environmental Protection: Subtitle F: Public Water Supplies: Chapter I: Pollution Control Board. Part 620: Groundwater Quality. Available at: <https://pcb.illinois.gov/SLR/IPCBandIPEAEnvironmentalRegulationsTitle35>

U.S. EPA (United State Environmental Protection Agency) Office of Solid Waste and Emergency Response (OSWER). 2013. Tier 3 Toxicity Value White Paper. OSWER Document Number 9285.7-86. Available at: <https://www.epa.gov/sites/production/files/2015-11/documents/tier3-toxicityvalue-whitepaper.pdf>