



ILLINOIS ENVIRONMENTAL PROTECTION AGENCY

2520 WEST ILES AVENUE, P.O. BOX 19276, SPRINGFIELD, ILLINOIS 62794-9276 • (217) 782-3397

JB PRITZKER, GOVERNOR

JAMES JENNINGS, ACTING DIRECTOR

HEALTH ADVISORY UPDATE FOR PERFLUOROBUTANOIC ACID (PFBA) CHEMICAL ABSTRACT SERVICES REGISTRY NUMBER (CASRN) 375-22-4

Prepared by:
Office of Toxicity Assessment
Illinois Environmental Protection Agency
April 11, 2025

REASON FOR ACTION

A new methodology to calculate health advisory guidance levels is now in effect, which lowers the guidance level for Perfluorobutanoic Acid (PFBA) from 0.007 milligrams per liter (mg/L), or 7,000 nanograms per liter (ng/L) or parts per trillion (ppt) to 0.0038 milligrams per liter (mg/L), or 3,800 nanograms per liter (ng/L) or parts per trillion (ppt).

On September 16, 2024, Illinois Environmental Protection Agency (Illinois EPA) issued a health advisory for Perfluorobutanoic Acid (PFBA) after confirming detection of the chemical in a well of a Community Water Supply (CWS). This sample result came from Illinois EPA's sampling investigation of Per- and Polyfluoroalkyl Substances (PFAS) in CWS statewide. 35 Illinois Administrative Code 620.605(a) directs the Illinois EPA to issue a health advisory for a chemical substance if all of the following conditions are met:

- 1) A community water supply well is sampled and a substance is detected and confirmed by resampling;
- 2) There is no standard under Section 620.410 for such chemical substance; and
- 3) The chemical substance is toxic or harmful to human health according to the procedures of Appendix A, B, or C.

The health advisory guidance level listed in the September 16, 2024 health advisory for PFBA was 0.007 milligrams per liter (mg/L), or 7,000 nanograms per liter (ng/L) or parts per trillion (ppt). The health advisory guidance level was calculated as required by the regulations at that time.

2125 S. First Street, Champaign, IL 61820 (217) 278-5800
115 S. LaSalle Street, Suite 2203, Chicago, IL 60603
1101 Eastport Plaza Dr., Suite 100, Collinsville, IL 62234 (618) 346-5120
9511 Harrison Street, Des Plaines, IL 60016 (847) 294-4000

595 S. State Street, Elgin, IL 60123 (847) 608-3131
2309 W. Main Street, Suite 116, Marion, IL 62959 (618) 993-7200
412 SW Washington Street, Suite D, Peoria, IL 61602 (309) 671-3022
4302 N. Main Street, Rockford, IL 61103 (815) 987-7760

On March 20, 2025, the Illinois Pollution Control Board adopted final amendments to Part 620, and on April 11, 2025, the amendments were published in the *Illinois Register*. These amendments update certain exposure factors used for calculating health advisory guidance levels, resulting in Illinois EPA's issuance of an updated PFBA health advisory level of 0.0038 milligrams per liter (mg/L), or 3,800 nanograms per liter (ng/L) or parts per trillion (ppt). The updated health advisory will be published in the Environmental Register (publication of the Illinois Pollution Control Board), and placed at the website: <https://pcb.illinois.gov/Resources/News>

The updated health advisory will also be placed on Illinois EPA's website at: <https://epa.illinois.gov/topics/water-quality/pfas/pfas-healthadvisory.html>

PURPOSE OF A HEALTH ADVISORY

In accordance with 35 Ill. Adm. Code 620.601, the purpose of a health advisory is to provide guidance levels that, in the absence of an applicable groundwater quality standard under Section 620.410, must be considered by Illinois EPA in: 1) establishing groundwater cleanup or action levels whenever there is a release or substantial threat of a release of a hazardous substance, pesticide, or another contaminant that represents a significant hazard to public health or the environment; 2) determining whether a community water supply is taking its raw water from a site or source consistent with regulatory requirements; and 3) developing Illinois Pollution Control Board (Board) rulemaking proposals for new or revised numerical standards.

Health advisories serve as informal technical guidance, intended to provide information about contaminant exposures and potential public health impacts. The guidance levels represent concentrations in drinking water at which no adverse health effects are expected to occur. Guidance levels are not enforceable or intended to be used as drinking water standards, also known as maximum contaminant levels (MCLs).

HEALTH ADVISORY GUIDANCE LEVEL FOR PFBA

Through issuance of this Health Advisory, Illinois EPA is providing public notice of its updated guidance level for PFBA in drinking water. For non-carcinogenic health effects, the updated guidance level is 0.0038 milligrams per liter (mg/L), or 3,800 nanograms per liter (ng/L) or parts per trillion (ppt).

Section 620.605 prescribes the methods for developing health advisories for carcinogens and non-carcinogens. PFBA does not meet the definition of a "carcinogen", as defined in Section 620.110; therefore, the method for developing a health advisory for non-carcinogens was used. Briefly, this method specifies that the United States Environmental Protection Agency (U.S. EPA) MCL or maximum contaminant level goal (MCLG) is the guidance level, if available, or the human threshold toxicant advisory concentration (HTTAC) must be determined using the procedures contained in Appendix A of Section 620. U.S. EPA has not published an MCL or

MCLG for PFBA; therefore, Illinois EPA used the Appendix A procedures to calculate a HTTAC for PFBA.

Appendix A specifies, in prescribed order, the toxicological data to be used in developing guidance levels. To determine appropriate toxicological data in accordance with nationally accepted guidelines, pursuant to the Illinois Groundwater Protection Act (415 ILCS 55-8(a)), Illinois EPA relied upon U.S. EPA guidance titled, “*Tier 3 Toxicity Value White Paper*” (paper), dated May 16, 2013, prepared by the U.S. EPA Office of Solid Waste and Emergency Response (OSWER) Human Health Regional Risk Assessors Forum. The paper lists a hierarchy of sources to be used when determining an appropriate toxicological value for use in human health assessments. The hierarchy for selection of toxicity values is as follows:

- Tier 1: U.S. EPA Integrated Risk Information System (IRIS).
- Tier 2: U.S. EPA Provisional Peer-Reviewed Toxicity Values (PPRTVs).
- Tier 3: In the order in which they are presented:
 - 1) The U.S. Health and Human Services Agency for Toxic Substances and Disease Registry (ATSDR) Dose Minimal Risk Levels (dose MRLs).
 - 2) California EPA, Office of Environmental Health Hazard Assessment (OEHHA).
 - 3) PPRTV “Appendix” Values.
 - 4) Health Effects Assessment Summary Table (HEAST).

The paper also references peer-reviewed toxicity values developed by other federal programs to calculate provisional drinking water health advisory levels as a Tier 3 source. In 2022, U.S. EPA placed Office of Water PFAS toxicity values above California EPA’s OEHHA toxicity values within the Tier 3 hierarchy.

In December 2022, U.S. EPA’s Integrated Risk Information System (IRIS) published a peer-reviewed toxicological assessment titled, “*IRIS Toxicological Review of Perfluorobutanoic Acid (PFBA, CASRN 375-22-4) and Related Salts.*” U.S. EPA’s IRIS toxicological assessment recommends a chronic oral reference dose (RfD) equal to 0.001 (1E-03) mg/kg-day. The value is based on a critical effect of increased liver weight and adverse thyroid effects in adult male rats from a study by Butenhoff et al. titled “*Toxicological evaluation of ammonium perfluorobutyrate in rats: twenty-eight day and ninety-day oral gavage studies,*” published in 2012. A no-observed-adverse-effect level (NOAEL) of 6 mg/kg-day was identified for NH₄⁺PFB, an ammonium salt of PFBA, and used to find the point of departure (POD) for PFBA by multiplying the NOAEL by the ratio of molecular weights (0.926) for a POD of 5.56 mg/kg-day. A human equivalent dose POD (POD_{HED}) of 1.27 mg/kg-day was then derived for oral PFBA exposure.

A total composite uncertainty factor (UF) of 1,000 (UF of 3 to account for toxicodynamic differences between humans and animals, UF of 10 to account for intraspecies variability, UF of 10 to account for extrapolation from subchronic to chronic, and UF of 3 to account for database uncertainties) was applied to the POD_{HED} .

The overall RfD for PFBA was calculated by dividing the POD_{HED} by the composite uncertainty factor.

$$RfD = \frac{POD_{HED}}{UF}$$

$$RfD = \frac{1.27 \text{ mg/kg-day}}{1,000}$$

$$RfD = 0.00127 \text{ mg/kg-day}$$

Rounded to one significant digit:

$$RfD = 0.001 \text{ mg/kg-day}$$


Using the RfD of 0.001 (1E-3) mg/kg-day, and the procedures outlined in Section 620. Appendix A, the recommended guidance level for drinking water is 0.0038 milligrams per liter (mg/L), or 3,800 nanograms per liter (ng/L) or parts per trillion (ppt).

CHEMICAL CHARACTERISTICS AND POTENTIAL ADVERSE HEALTH EFFECTS

General Description of PFBA

Perfluorobutanoic Acid (CASRN 375-22-4), also known as heptafluorobutyric acid or PFBA, is a synthetic chemical which is part of a larger class of chemicals referred to as per- and polyfluoroalkyl substances. PFAS have been manufactured since the middle 20th Century and are known for their chemical and physical properties that impart oil and water repellency, temperature resistance, and friction reduction to a wide range of products, including, but not limited to, textile coatings, paper products, food wrappers, cosmetic and personal care products, non-stick cookware and fire-fighting foams. PFAS are also used in the semiconductor, aerospace, oil production and mining, and metal plating industries, to name a few. PFAS enter the environment through industrial manufacturing and the use and disposal of PFAS-containing products. The chemical and physical properties of PFBA make it mobile, persistent and bioaccumulative, meaning fish and other animals may accumulate PFBA in animal tissue when their food sources are contaminated with PFBA. PFBA is known to be persistent in the environment.

Structural Identifier



Chemical Identifier



Potential Adverse Health Effects of PFBA

Epidemiological studies on human health effects from exposure to PFBA are limited in their ability to draw conclusions on the associations between health effects and exposure.

Information regarding health effects of PFBA are primarily derived from animal studies, via the ingestion, or oral exposure, route. Laboratory studies observed the following effects in animals exposed to PFBA:

- Increased relative liver weight
- Increased hepatocyte hypertrophy
- Increased thyroid hormone T4
- Embryo/fetal mortality
- Developmental delays

Carcinogenic Potential

Section 620.110. defines a carcinogen as a contaminant that is classified as: 1) a Category A1 or A2 Carcinogen by the American Conference of Governmental Industrial Hygienists (ACGIH); 2) a Category 1 or 2A/2B Carcinogen by the World Health Organization's International Agency for Research on Cancer (IARC); 3) a "Human Carcinogen" or "Anticipated Human Carcinogen" by the United States Department of Health and Human Service National Toxicological Program (NTP); or 4) a Category A or B1/B2 Carcinogen or as "carcinogenic to humans" or "likely to become carcinogenic to humans" by the U.S. EPA in IRIS or a Final Rule issued in a Federal Register notice by the USEPA. PFBA is not classified as a carcinogen by any of the above sources.

**ATTACHMENT TO HEALTH ADVISORY
FOR
PERFLUOROBUTANOIC ACID (PFBA)
CASRN 375-22-4**

OVERVIEW OF KEY STUDIES

For information regarding the studies used by U.S. EPA's IRIS for the derivation of its PFBA RfD, refer to *IRIS Toxicological Review of Perfluorobutanoic Acid (PFBA, CASRN 375-22-4) and Related Salts*, located at:

https://cfpub.epa.gov/ncea/iris_drafts/recordisplay.cfm?deid=350051.

DERIVATION OF THE HEALTH ADVISORY GUIDANCE LEVEL FOR PFBA

The first step in the derivation of a health advisory guidance level is to determine whether the chemical substance presents a carcinogenic risk to humans. PFBA does not meet the definition of a carcinogen as specified in Part 620. Therefore, the guidance level will be based on non-carcinogenic effects of this chemical.

In deriving a guidance level to protect against a health effect for which there is a threshold dose below which no damage occurs (i.e., noncarcinogen effects), Section 620.605 specifies that U.S. EPA's MCLG, if available, is the guidance level. U.S. EPA has not published a MCLG for PFBA; therefore, Illinois EPA must calculate the HTTAC as the guidance level, using the procedures specified in Appendix A of Section 620.

Appendix A specifies in subsection (a) that the HTTAC is calculated as follows:

$$HTTAC = \frac{RSC \cdot ADE}{W}$$

Where:

HTTAC = Human threshold toxicant advisory concentration in milligrams per liter (mg/L).

RSC = Relative source contribution, the relative contribution of the amount of exposure to a chemical via ingestion of drinking water when compared to total exposure to that chemical from all sources. Valid chemical-specific data must be used if available. If valid chemical-specific data are not available, a value of 20% (= 0.20) must be used.

ADE = Acceptable daily exposure of a chemical in milligrams per day (mg/d) as determined under Appendix A, subsection (b).

W = Per capita daily water consumption for a child (0-6 years of age), equal to 0.78 liters per day (L/d).

Subsection (b) of Appendix A specifies that the ADE be calculated using, in specified order: a U.S. EPA verified RfD (an estimate of a daily exposure to a chemical which is expected to be without adverse health effects for humans for a lifetime of exposure in units of mg/kg-day); the most sensitive Point of Departure (POD) as determined by Benchmark Dose Modeling or the NOAEL/LOAEL approach consistent with U.S. EPA RfD guidance; and derivation of a Human Equivalent Dose (HED) using physiologically based pharmacokinetic (PBPK) modeling or Dose Adjustment Factor (DAF), then divided by the total Uncertainty Factor (UF) and modifying factor (MF), if applicable. .

Illinois EPA selected the U.S. EPA IRIS recommended RfD of 0.001 (1E-3) mg/kg-day, as the verified RfD for use in calculating the ADE. The ADE equals the product of multiplying the toxicity value by 15 kilograms (kg), which is the updated assumed average body weight of a child age 0-6 years per Section 620:

$$ADE = 0.001 \text{ mg/kg-day} \cdot 15 \text{ kg} = 0.015 \text{ mg/day}$$

The next step in the development of the HTTAC is the evaluation of chemical-specific RSC data available for the chemical. Illinois EPA evaluated data from ATSDR, U.S. EPA Office of Water, and values developed by other states. There is little scientific consensus regarding the contribution of drinking water to the total amount of PFAS exposure to humans. Humans are exposed to PFBA through a variety of media, including, but not limited to air emissions, ingestion of fish or other animals exposed to PFBA, dermal exposure and incidental exposure from PFBA-containing consumer products, much of which varies on a site-specific basis. Due to this lack of consensus, Illinois EPA elected to use the conservative default value of 20% (0.20) for its HTTAC calculation.

The HTTAC is calculated by the product of the RSC and the ADE, divided by the per capita daily water ingestion rate for a child (0-6 years of age), specified in Appendix A as equal to 0.78 L/day:

$$HTTAC \text{ (mg/L)} = \frac{0.20 \cdot 0.015 \text{ mg/day}}{0.78 \text{ L/day}}$$

$$HTTAC \text{ (mg/L)} = \frac{0.003 \text{ mg/day}}{0.78 \text{ L/day}}$$

$$HTTAC = 0.0038 \text{ mg/L}$$

or:

$$3,800 \text{ ng/L or ppt}$$

The final step in ensuring a calculated guidance level is appropriate is to compare the guidance level to the chemical's practical quantitation limit (PQL), or minimum reporting level (MRL). U.S. EPA's Method 537.1 for analyses of PFAS drinking water samples shows the PFBA MRL is 1.8 ng/L, which is below the calculated guidance level of 3,800 ng/L. Therefore, the guidance level is appropriate.

REFERENCES

Butenhoff, JL; Bjork, JA; Chang, SC; Ehresman, DJ; Parker, GA; Das, K; Lau, C; Lieder, PH; van Otterdijk, FM; Wallace, KB. (2012). Toxicological evaluation of ammonium perfluorobutyrate in rats: twenty-eight-day and ninety-day oral gavage studies. *Reproductive Toxicology*, vol. 33. 513-530. Available at:

<http://www.sciencedirect.com/science/article/pii/S0890623811003522>

IGA (Illinois General Assembly). Illinois Groundwater Protection Act (IGPA). 415 ILCS 55.

Available at: <https://www.ilga.gov/legislation/ilcs/ilcs3.asp?ActID=1595&ChapterID=36>

PCB (Pollution Control Board). Title 35: Environmental Protection: Subtitle F: Public Water Supplies: Chapter I: Pollution Control Board. Part 620: Groundwater Quality. Available at:

<https://pcb.illinois.gov/SLR/IPCBandIEPAEnvironmentalRegulationsTitle35>

U.S. EPA (United State Environmental Protection Agency) Office of Solid Waste and Emergency Response (OSWER). 2013. Tier 3 Toxicity Value White Paper. OSWER Document Number 9285.7-86. Available at: <https://www.epa.gov/sites/production/files/2015-11/documents/tier3-toxicityvalue-whitepaper.pdf>

U.S. EPA IRIS Toxicological Review of Perfluorobutanoic Acid (PFBA, CASRN 375-22-4) and Related Salts. U.S. Environmental Protection Agency, Washington, DC, EPA/635/R-22/277Fb, 2022. Available at: https://cfpub.epa.gov/ncea/iris_drafts/recordisplay.cfm?deid=350051